



E-ISSN: 2663-1067  
P-ISSN: 2663-1075  
IJHFS 2022; 4(2): 39-42  
Received: 12-05-2022  
Accepted: 18-07-2022

**Bankar PN**  
Department of Horticulture,  
Vasant Rao Naik Marathwada  
Krishi Vidyapeeth, Parbhani,  
Maharashtra, India

**Mane RR**  
Department of Horticulture,  
AKS, university Satana,  
Madhya Pradesh, India

**Dr. Tambe TB**  
Department of Horticulture,  
Vasant Rao Naik Marathwada  
Krishi Vidyapeeth, Parbhani,  
Maharashtra, India

**Corresponding Author:**  
**Bankar PN**  
Department of Horticulture,  
Vasant Rao Naik Marathwada  
Krishi Vidyapeeth, Parbhani,  
Maharashtra, India

## Effect of levels of nitrogen with boron and zinc on growth of broccoli (*Brassica oleraceae* L. Var. *italica*) Cv. Phule Ganesh

**Bankar PN, Mane RR and Dr. Tambe TB**

### Abstract

The investigation was carried out in shadenet of Instructional- cum- Research farm of Department of Horticulture, College of Horticulture, VNMKV, Parbhani during the year 2018-2019. The seven different treatments including control were used in Randomized Block Design (RBD). There were significant differences found in growth, yield and quality by the soil application of NPK and B and Zn with different treatments. Application of 130 kg N + 60 kg P<sub>2</sub>O<sub>5</sub> + 15 kg B ha<sup>-1</sup> gave maximum plant height (29.57 cm), stem diameter (4.06 cm), Number of leaves (20.63), Length of longest leaves (26.37 cm), Width of leaves (16.70 cm), leaf area (717.07 cm<sup>2</sup>), leaf area index (0.199), days taken to central curd formation (53.43), days taken to central curd maturity (72.77), curd diameter (16.18 cm), curd length (14.20 cm), weight of central curd per plant (326.73 g).

**Keywords:** Broccoli, NPK, boron, and zinc

### 1. Introduction

Broccoli (*Brassica oleracea* L. var. *italica*) Cv. Phule Ganesh” belongs to the genus *Brassica*, and family Brassicaceae which includes a wide range of crop plants derived from the Mediterranean sea and modified over the years by selection and breeding (Decoteau, 2000) [3]. Vegetables defined as those herbaceous plants of which some portion is eaten either cooked or raw, during the principal part of the meal, constitute an important segment of our agricultural system. Vegetables are rich source of carbohydrates, proteins, vitamins and minerals. India stands at second place in vegetable production after China. (Thamburaj and Singh, 2003) [11]. China is the top world producer of broccoli (*Brassica oleracea* L. var. *italica*) followed by India, USA, Spain, Italy, France, Mexico, Poland, Pakistan and United Kingdom (FAO, Statistics 2018) [4]. In India broccoli is under-utilized food crop, mainly confined to limited area especially near the big cities. In India annual production around 180684 million tonnes of vegetable from 10172 million hectares of land (Anonymous, 2018) [2]. However, this production does not meet the requirement of 300 g of vegetables per capita per day. Thus, with the large vegetarian population, the production of vegetables in India needs to be greatly increased. Though the central and state governments have taken some steps for increasing the production of vegetable crops, it needs further boost from the research activities.

Exotic vegetables are increasingly popular in star hotels. These vegetables generally require cool temperature. The agro-climatic conditions prevailing in several part of northern Himalayas of Jammu and Kashmir, Himachal Pradesh and Kumaon and Garhwal division of Uttaranchal are very favorable for growing these vegetables. Besides, they may also be grown during winter season in plains of North India (Pandey and Rai, 2005) [7]. The family Brassicaceae is an important for horticulture because there are several commercial vegetable crops like cabbage, cauliflower, Brussels sprouts, sprouting broccoli, etc. which are cultivated extensively; further the members of the Brassicaceae kind are characterized by the fact that they contain sulphur which is an important constituent of a balanced human diet. Broccoli is cool season crop. The temperature of 20 °C to 25 °C is optimum for its proper growth while 15 °C to 20 °C for its heading stage. It is more sensitive to temperature. When the plants are small and tender, they are susceptible to cold injury. Warm weather is disadvantageous, since the bud clusters grow loose quickly. In Northern India, generally, it is planted in September and October and is ready for harvest from late November to early December and may continue till early February.

Broccoli needs fertile soil. Medium black soil is suitable for broccoli cultivation. Soil pH ranges from 6.5 to 7.0. In this crop, planting distance needs to be maintained properly to avoid pest problem and to get quality curds for the market. The distance can be reduced to 45x45 cm and 60x60 to avoid stem hollowness in the stem. Broccoli is an important vegetable crop and has high nutritional and good commercial value (Yoldas, *et al.* 2008) [13]. Broccoli is low in sodium food, fat free and calories, high in vitamin C and good source of vitamin A, B1, B2 and calcium (Decoteau, 2000) [3]. Now-a-days, broccoli attracted more attention due to its multifarious use and great nutritional value (Rangkadilok, *et al.* 2004) [8]. Broccoli is a rich source of vitamins, minerals, proteins, etc. It has about 130 times more vitamin-A contents than cauliflower and 22 times more than cabbage. It is also a rich source of sulphoraphane, a compound associated with reducing risk of cancer. It contains vitamin A (9000 mg-100 g), vitamin B (33 mg-100 g), vitamin C (137 mg-100 g), minerals *viz*; Ca (1.29%), P (0.79%), K (3.5%), S (1.26%), Fe (205 ppm), I (1.965 ppm), Cu (24 ppm), protein (3.3%), total carbohydrates (5.5%), fat (0.2%), water (89.9%) and calories (36-100 g) (Thamburaj and Singh, 2003) [11]. The Marathwada region is totally new area for its cultivation. However, it is need to produce better quality curds. Micronutrients like boron and zinc play the important role to improve growth, yield and quality of broccoli. Beside the crop duration and the harvesting duration, sprouting broccoli is shorter than cabbage and cauliflower. The response may differ in growth, yield and quality parameters of the broccoli. In view of above present.

## 2. Materials and Methods

An experiment was conducted at College of Horticulture, Parbhani. Geographically Parbhani is situated at 19° 16' North latitude and 76° 47' East longitude and at 408.50 meter above sea level in Marathwada division encompassed by 17° 35' to 24° 40' North latitude and 74° 49' to 78° 15' East longitude geographical boundaries. The meteorological data for corresponding period of crop season recorded at central meteorological observatory, Vasantrao Naik Marathwada Krishi Vidyapeeth, Parbhani. The perusal of data indicated that the average maximum and minimum temperatures during crop growing period were 33.04 °C and 14.8 °C, respectively. The mean relative humidity (RH) of morning (RH I) and evening (RH II) hours were 78.6 and 40.5 per cent. The mean sunshine hours per day were 8.6 hours day<sup>-1</sup>. The mean wind velocity was 4.3 km h<sup>-1</sup>.

The total precipitation received during crop growth period in 2017-18 was 21.54 mm. The evaporation was 4.72 mm recorded during the crop growing season. The present experiment entitled "Effect of levels of nitrogen with boron and zinc on growth and yield of broccoli. (*Brassica oleracea* L. var. *italica*) Cv. Phule Ganesh". The field experiment was carried out in Randomized Block Design (RBD) with three replications and seven treatments *viz*; T1 - 120:60:40 NPK kg-ha with boron at 10 kg-ha, T2 - 130:60:40 NPK. Investigation is entitled "Effect of levels of nitrogen with boron and zinc on growth, yield and quality of broccoli (*Brassica oleracea* L. var. *italica*) Cv. Phule Ganesh. Kg-ha with boron at 15 kg-ha, T3 - 140:60:40 NPK kg-ha with boron at 20 kg-ha, T4 - 120:60:40 NPK kg-ha with zinc at 10 kg-ha, T5 - 130:60:40 NPK kg-ha with zinc at 15 kg-ha, T6 - 140:60:40 NPK kg-ha with zinc at 20 kg-ha and T7 - control. The results

obtained during the course of investigation are summarized below. The seeds of earlier mentioned variety of broccoli were sown in First week of October in portraits filled with cocopeat. Thereafter, 30 days old seedlings of broccoli were transplanted in well prepared experimental field at distance 60 X 60 cm. At the time of transplanting, half dose of nitrogen, full dose of Phosphorous and potash were applied in experimental plots and boron and zinc later apply with 15 days after transplanting, remaining half dose of nitrogen was applied after one month of transplanted crop. The plant protection measures were also followed to control of pest and diseases. To find out the effect of treatments, five sample plants were selected to obtain field data according to observation on growth and yield of broccoli.

## 3. Result and Discussion

**3.1 Effect of levels of NPK with boron and zinc on growth parameters data exhibited from the table 1.** Showed that the application of nitrogen, phosphorous, potash and boron had significant effect on the plant height of broccoli Cv. Phule Ganesh. Each increment of NPK, boron and zinc doses upto 130:60:40 kg/ha NPK + Boron 15 kg/ha were obtained significantly height of plant 60 DAT (29.27 cm), The maximum stem diameter at 30 DAT (3.06 cm) and at 60 DAT (4.06 cm) was recorded by the application of T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha. The maximum number of leaves per plant was recorded at 30 DAT (15.10) and at 60 DAT (20.63) was recorded by the treatment T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha. The maximum length of leaves of plant recorded at 30 DAT (23.68 cm) and at 60 DAT (26.37 cm) was recorded by the treatment T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha. The maximum width of leaves of plant was recorded at 30 DAT (14.07cm) and at 60 DAT (16.70 cm) by the treatment T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha. The maximum leaf area of broccoli (717.07cm<sup>2</sup>) was recorded by the treatment T2 *i.e.* 130:60:40 NPK Kg-ha with boron at 15 Kg-ha. The maximum leaf area index of broccoli (0.199) was recorded by the treatment T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha. The minimum days required for curd formation DAT of broccoli (53.43) was recorded by the treatment T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha. The minimum days required for curd maturity DAT of broccoli (72.77) was recorded by the treatment T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha maximum curd diameter of broccoli (16.18 cm) was recorded by the treatment T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha. The maximum curd length of broccoli (14.20 cm) was recorded by the treatment T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha. The maximum weight of curd of broccoli (326.73 g) was recorded by the treatment T2 *i.e.* 130:60:40 NPK kg-ha with boron at 15 kg-ha. The application of boron after transplanting was found significant for maximum height of plant in broccoli. stem diameter, Number of leaves, Length of longest leaves, Width of leaves, leaf area, leaf area index, days taken to central curd formation, days taken to central curd maturity, curd diameter, curd length, weight of central curd per plant This might be due to the stimulating influence of boron enhancing the rate of absorption of N, P and K and other nutrients. Moniruzzaman *et al.* (2007) [14], Moreover, boron took part in sugar translocation which might be lead to the increased height of plant. This is in accordance with the findings of similar results observed Saha *et al.* (2010) [9], Naher *et al.* (2014) [6] on cabbage. Singh *et al.* (2015) [10] on

broccoli. The plant which treated with nitrogen (N) grows quickly and retain maximum vegetative growth by means of higher canopy area development, CO<sub>2</sub> exchange rate and photosynthetic activity Yasir *et al.* (2016) [12] potassium gives role on carbohydrate metabolism and enzyme

activation in plant body and boron act as a proper translocation of sugars, starch and nitrogen compound in plant body these finding are in close conformity with earlier results obtained by Alam *et al.* (2007) [1] and Singh *et al.* (2015) [10].

**Table 1:** Effect of levels of NPK with boron and zinc on growth and leaf attributes at 30 days

Sr. No.	Treatment details	Plant height (cm)	Stem diameter (cm)	Number of leaves	Length of Longest leaf (cm)	Width of leaves (cm)
T1	120:60:40 NPK kg-ha+ B 10 kg-ha	18.93	2.43	14.20	21.83	12.46
T2	130:60:40 NPK kg-ha+ B 15 kg-ha	22.60	3.06	15.10	23.68	14.07
T3	-ha+ B 20 kg-ha	18.20	2.2	13.40	21.00	12.30
T4	120:60:40 NPK kg-ha+ Zn 10 kg-ha	18.83	2.3	12.86	21.40	12.63
T5	130:60:40 NPK kg-ha+ Zn 15 kg-ha	21.83	2.53	14.33	23.00	13.03
T6	140:60:40 NPK kg-ha+ Zn 20 kg-ha	18.00	2.03	12.80	20.40	12.40
T7	Control	16.60	1.8	11.40	20.03	11.50
	S.E.(±)	0.67	0.10	0.42	0.68	0.39
	C. D. 5%	2.06	0.29	1.28	2.10	1.21

**Table 2:** Effect of levels of NPK with boron and zinc on growth and leaf attributes at 60 days

Sr. No.	Treatment details	Plant height (cm)	Stem diameter (cm)	Number of leaves	Length of Longest leaf (cm)	Width of Leaves (cm)
T1	120:60:40 NPK kg-ha +B 10 kg-ha	25.67	3.50	19.07	25.17	14.60
T2	130:60:40 NPK kg-ha + B 15 kg-ha	29.27	4.06	20.63	26.37	16.70
T3	140:60:40 NPK kg-ha + B 20 kg-ha	24.70	2.96	16.07	24.60	14.47
T4	120:60:40 NPK kg-ha +Zn 10 kg-ha	25.27	3.43	17.80	24.63	14.93
T5	130:60:40 NPK kg-ha + Zn 15 kg-ha	28.03	3.68	20.03	25.73	15.43
T6	140:60:40 NPK kg-ha + Zn 20 kg-ha	23.87	3.16	15.27	23.57	13.73
T7	Control	20.03	2.4	14.80	21.93	12.60
	S.E.(±)	0.80	0.17	0.52	0.62	0.60
	C. D. 5%	2.48	0.54	1.60	1.91	1.91

**Table 3:** Effect of levels of NPK with boron and zinc on leaf attributes

Sr. No.	Treatment details	Leaf area (cm <sup>2</sup> )	Leaf area index
T1	120:60:40 NPK kg <sup>-ha</sup> + B 10 kg <sup>-ha</sup>	18.93	2.43
T2	130:60:40 NPK kg <sup>-ha</sup> + B 15 kg <sup>-ha</sup>	22.60	3.06
T3	140:60:40 NPK kg <sup>-ha</sup> + B 20 kg <sup>-ha</sup>	18.20	2.2
T4	120:60:40 NPK kg <sup>-ha</sup> + Zn 10 kg <sup>-ha</sup>	18.83	2.3
T5	130:60:40 NPK kg-ha + Zn 15 kg-ha	21.83	2.53
T6	140:60:40 NPK kg-ha + Zn 20 kg-ha	18.00	2.03
T7	Control	16.60	1.8
	S.E.(±)	0.67	0.10
	C. D. 5%	2.06	0.29

**Table 4:** Effect of levels of NPK with boron and zinc on curd attributes

Sr. No.	Treatment details	Days taken To central curd formation	Days Taken to central curd maturity	Curd diameter (cm)	Curd length (cm)	Weight of Curd per plant (g)
T1	120:60:40 NPK kg-ha+ B 10 kg-ha	56.23	74.30	14.63	12.50	277.07
T2	130:60:40 NPK kg-ha+ B 15 kg-ha	53.43	72.77	16.18	14.20	326.73
T3	140:60:40 NPK kg-ha+ B 20 kg-ha	60.20	79.20	13.56	11.83	275.53
T4	120:60:40 NPK kg-ha+ Zn 10 kg-ha	55.70	77.53	14.60	12.76	294.37
T5	130:60:40 NPK kg-ha+ Zn 15 kg-ha	54.13	73.13	15.09	13.40	310.4
T6	140:60:40 NPK kg-ha+ Zn 20 kg-ha	62.90	80.83	13.24	11.76	282.23
T7	Control	63.60	84.90	12.36	11.60	211.6.
	S.E.(±)	1.14	1.67	0.55	0.48	5.48
	C. D. 5%	3.52	5.14	1.70	1.47	16.45

## Conclusion

The present findings have clearly indicated that the combined soil application of different nitrogen levels and micronutrient boron and zinc at 130:60:40 NPK Kg-ha + boron at 15Kg-ha and 130:60:40 NPK Kg-ha + zinc at 15 Kg-ha 15 DAT may be suggested as the optimum level for improving growth broccoli.

## References

1. Alam MN. Effect of Boron Levels on Growth and Yield of Cabbage in Calcareous Soils of Bangladesh. Res. J of Agril. And Bio. Sci. 2007;3(6):858-865.
2. Anonymus. Indian Horticultural Database, National Horticultural Board, Government of India; c2018.
3. Decoteau DR. Vegetable Crops. Prentice Hall, Upper

- Saddle River, New Jersey, ISBN. 2000;10(3):464.
4. FAO Food and Agricultural Organization of the United Nations. Cauliflower and Broccoli Area and Production data; c2018. <http://www.fao stat.org.com>.
  5. M Rahman SM, Kibria MG. Effect of boron and nitrogen on yield and hollowstem of broccoli. J sci. nature. 2007;1(3):24-29.
  6. Naher MN, Alam MN, Jahan N. Effect of Nutrient Management on the Growth and Yield of Cabbage (*Brassica oleracea* var. *capitata* L.) in Calcareous Soils of Bangladesh. The Agriculturists. 2014;12(2):24-33.
  7. Pandey AK, Rai M. Popularizing exotic vegetables for nutrition and food diversification. Indian Hort. 2005.Jul-Sep:19-22 and 36.
  8. Rangkadilok N, Nicolas ME, Bennet RN, Eagling DR. The effect of sulphur fertilizer on glucoraphanin levels in broccoli (*Brassica oleraceae*. L. var *italica*) at different growth stages. J of agril. and food chem. 2004;52:2632-2639.
  9. Saha P, Ranidas N, Chatterjee R. Boron and molybdenum nutrition IN sprouting broccoli under terai of West Bengal. As. J of Hort. 2010;5(2):353-355.
  10. Singh MK, Chand T, Singh KV. Responses of different doses of NPK and boron on growth and yield of broccoli. Int.J of Bio-resource and stress management. 2015;6(1):108-112.
  11. Thamburaj S, Singh N. Textbook of Vegetables, Tuber Crops and Spices: ICAR New Delhi; c2003. p. 136-137.
  12. Yasir EM, Ahmed AM, Osman A, Makeen, Tarig EA Suliman TE. Effect of NPK micro doses fertilizer on leaf area, leaf area index and pods and hay yield of six genotypes of groundnut-north kordofan state-sudan Int. J OF Sci. & Tech. Res. 2016;5(5):2277-8616.
  13. Yoldas F, Ceylan S. Effect of nitrogen fertilizer on yield, quality and nutrient content broccoli. J of pl. nutrition. 2008;31:1333-1343.
  14. Moniruzzaman M, Chattopadhyay J, Billups WE, Winey KI. Tuning the mechanical properties of SWNT/Nylon 6, 10 composites with flexible spacers at the interface. Nano letters. 2007 May 9;7(5):1178-85.