



E-ISSN: 2663-1067
P-ISSN: 2663-1075
<https://www.hortijournal.com>
IJHFS 2023; 5(1): 91-95
Received: 18-03-2023
Accepted: 25-04-2023

Rajdeep Singh
Assistant Professor,
Department of Agriculture,
Rayat Bahara, Hoshiarpur,
Punjab, India

Jasmine
Assistant Professor,
Department of Agriculture and
Botany, GKSM Government
College, Tanda Urmar,
Hoshiarpur, Punjab, India

Dr. Sonia
Assistant Professor,
Department of Agriculture and
Botany, GKSM Government
College, Tanda Urmar,
Hoshiarpur, Punjab, India

Corresponding Author:
Rajdeep Singh
Assistant Professor,
Department of Agriculture,
Rayat Bahara, Hoshiarpur,
Punjab, India

To evaluate the effect of different pre-chemical treatments and packing materials on the fruit quality of mosambi (*Citrus limetta*)

Rajdeep Singh, Jasmine and Dr. Sonia

DOI: <https://doi.org/10.33545/26631067.2023.v5.i1b.164>

Abstract

A study was carried out in entitled “To evaluate the effect of different pre-chemical treatments and packing materials on the fruit quality of Mosambi (*Citrus limetta*)” at the Agricultural laboratory of Sant Baba Bhag Singh University, Jalandhar, Punjab (India) during the year 2020. Completely Randomized Design (CRD) with 3 replications and 10 treatments was constructed for the present study. Fruits were pre-treated in hot water (55 °C) was given for 1 minute. Then, fruits were air dried and packed in packaging films viz. high-density polyethylene (HDPE) and other packaging materials, Low-density polyethylene (LDPE) with or without perforations) and sealed with an electronic sealing system after being dehumidified. Additionally, chemically viz. 2, 4-dichlorophenoxyacetic acid (2, 4-D) and gibberellic acid (GA3), prepared fruits were also kept at (ambient) room temperature. For analyzing, fruit samples were collected for various quality parameters after 7 days of interval of storage (0, 7th, 14th, and 21th). The results revealed that among the different treatments T₄ (HDPE perforated) treatment was superior for characters like phenol content, TSS, total titrable acid, total sugars, reducing sugars and non-reducing sugars, as compared to other treatments studied. On the other hand, fruit pH and ascorbic acid was observed better in T₆ and T₃ respectively. It was concluded that the use of HDPE perforated packaging material might be effective and economical treatment for improving the quality of Mosambi fruit.

Keywords: 2, 4-D, HDPE and CRD, economical

1. Introduction

Mosambi (*Citrus limetta*) belongs to Rutaceae family which is a self-pollinated fruit crop and can be grown successfully in both tropical and sub-tropical areas. *Citrus limetta* is the result of the cross between citron (*Citrus medica*) and bitter orange (*citrus × aurantium*) [2]. Citrus fruits are low in salt, fat, and cholesterol but high in minerals (potassium, calcium, phosphorus, magnesium, copper), vitamins (C, B1, B2, B3, B5, B6, B9), fiber, and phytochemicals (carotenoids, phenols, including flavonoids, coumarins, limonoids, alkaloids, and essential oils) all of which have been shown to reduce inflammation when taken alone or in combination with nervous system disorders, heart disease, and even cancer [13]. Citrus is one some of the widely grown fruit crops within the international with annual worldwide production of 102 MT approximately and in India region under citrus cultivation is 973 thousand hectares with annual production of 122 lakh MT during 2018-19. [7]. In harvested fruits, loss in water vapour, results in shrinkage with peel, turgidity reduction and lowered resistance to gas diffusion, results in negative consequences with the taste and flavour [4]. To reduces post-harvest losses and enhancing the keeping quality of citrus fruits, different post-harvest treatments play a vital role. Therefore, this current work was evaluated the effect of different pre-chemical treatments and packing materials on the fruit quality of Mosambi.

Methods and Materials

The experiment was conducted in Completely Randomized Design with three replications and 10 treatments for analyzing the effect of different pre-chemical treatments and packaging material on qualitative traits of Mosambi in the laboratory of Punjab agro. Juice ltd. Hoshiarpur FSSC LIC NO.10014063000386 and at Lal path lab, Nandachor in December, 2020.

The total number of treatments were analyzed for 7 days of interval (0, 7th, 14th, and 21th). Pretreatment of hot water (55 °C) was given for 1 minute. Then treated fruits were air dried and packed in packaging films *viz.* high-density polyethylene (HDPE) and other packaging materials, Low-density polyethylene (LDPE) with or without perforations) and sealed with an electronic sealing system after being dehumidified. Chemically *viz.* 2, 4-dichlorophenoxyacetic acid (2, 4-D) and gibberellic acid (GA3), prepared fruits were also kept at (ambient) room temperature. The fruits samples were chopped with stainless steel knife and sample was pulped with pestle and mortar using distilled water for analyzing for biochemical characteristics *viz.* Fruit pH, Phenol content, total soluble solids (°Brix), Total titratable acidity (%), total sugars, reducing sugars, non-reducing sugars and ascorbic acid. Therefore, present investigation was conducted with 10 different treatments *viz.*, T₁ (LDPE), T₂ (LDPE perforated), T₃ (HDPE), T₄ (HDPE perforated), T₅ (Cold storage), T₆ (Treated with 2,4-D 50 ppm (without

packaging)), T₇ (Treated with 2,4-D 100 ppm (without packaging)), T₈ (Treated with GA3 50 ppm (without packaging)), T₉ (Treated with GA3 100 ppm (without packaging)), T₀ (Control).

Results and Discussion

The analysis of variance revealed significant difference among all the treatments and traits under the study. Data evaluated on the effect of different pre-chemical treatments and packing materials on the fruit quality of Mosambi.

FRUIT pH: The minimum pH recorded in refrigerated storage (T₆) compared with all treatments such as for low-density polyethylene (T₁) packaging and for fruit pre-chemically processed 2, 4-D 100 ppm (T₇), was the minimal value observed presented in Fig. 1. The maximum pH all through the test was recorded in the control (T₀). Sushma also studied that fruit pH fluctuated all through storage ^[16].




Fig 1: Effect of different treatments on pH of fruit

Phenol content

The data from this phenol test showed that the phenol value increased with increasing storage time. The results showed that random and differential treatments had a significant effect on phenol levels in mosses on the last day or end of storage. In packaging materials, perforated high-density

polyethylene (T₄) has a phenol content value of at least 250 compared to other equipment. The maximum value of phenol turned into shown in control (T₀) 543 as shown in Fig 2. Ingle also said that end result which are treated with the 2, 4-D chemical were effective for reducing the phenol content in citrus ^[8].




Fig 2: Effect of various treatments on the phenol content of fruit

Total soluble solids (TSS): According to the results of this analysis, the total soluble solids (TSS) increase as the storage time increases, as shown in Fig 3. Significantly minimum value 10.2 °Brix of total soluble solid was recorded in high-density polyethylene with perforation (T₄) at the end day of storage. Whereas, the maximum value of TSS recorded in control (T₀). Rab *et al.* [14] said that rapid

growth in TSS at some stage in storage because of the decrease in acidity and this affected the TSS of the fruits. Saltviet [15] stated that fruits full of perforated high-density polyethylene at low temperature confirmed low TSS level due to the fact fruit packaging of decreased the respiratory and dehydration, it will greater effective while packed fruits stored in low temperature.




Fig 3: Effect of various treatments on TSS of fruit

Total titratble acid (%)

Data pertaining to treatment means showed that the utmost worth of TTA was recorded in perforated high density polythene (T₄). Minimum values were recorded within the control (T₀) compared to any or all treatments presented in Fig 4. Echeverria and Valich investigated that the decrease in TTA throughout storage might be caused by fruit

respiration through the employment of organic acids, resulting in quicker fruit ripening [5]. Vines and Ober Bacher said that [HDP] packaging is more effective or useful for holding the quantity of acid at low temperatures as a result of the changed atmosphere in polythene limits the respiration rate [18].




Fig 4: Effect of different treatments on the total Titrable acid

Total sugar

To maintain good fruit quality, it's necessary to prevent the expansion of total sugar. The data shown in Fig 5 indicates that the fruits treated with completely totally different treatments had a significant impact on the total sugar of mosambi fruit. The perforated high density polyethylene (T₄) packaging material has a minimum value (10.4%)

compared to any or all different treatments. The maximum (12.9%) recorded total sugar value was achieved in unprocessed fruit (T₀) compared to additional processed fruit. Keditu investigated that high sugar content was noted within the controls because of moisture loss and reaction of starch (polysaccharide) to sugar (monosaccharide) in citrus fruit (Kinnow mandarin) [11].




Fig 5: Effect of different treatments on the total sugar of fruit

Non-reducing sugar: The minimal value (4.2%) of non-reducing sugar was recorded in HDPE perforated (T₄) and maximum value (6.5%) was recorded in control (T₀) compared to other treatments showed in Fig 6. Ali supports the current results, stating that the drop in non-reducing sugar may be attributed to the conversion of non-reducing

sugar to reducing sugar [1]. Kaur also stated that rapid decrease in non-reducing sugar might to ambient conditions because temperature in ambient conditions remain high compared to recommended storage and advanced temperature lower the relative moisture results high transpiration rate or water loss [10].




Fig 6: Effect of different treatments on the non-reducing sugar of fruit

Reducing sugar: A small amount of sugar reduction (5.2%) was recorded on packaging made of high-density polyethylene with perforation (T₄) and a higher rate (6.4%) of sugar reduction was recorded in control (T₀) compared to all treatments shown in Fig 7. Jawanda studied the reason

behind the rapid increase in reducing sugar might be respiration which results the dehydration of juice content of fruit and conversion of polysaccharides into monosaccharide [9].




Fig 7: Effect of different treatments on the reducing sugar of fruit

Ascorbic ACID

Based on this study, it was investigated that ascorbic acid in the fruit decreased with late-term growth and all different treatments had a significant effect on the ascorbic acid content in Mosambi fruit presented in Fig 8. It was analyzed that high levels of ascorbic acid (45) was found in high-density polyethylene (T₃). However, the minimum value

(36) was recorded in control (T₀). The findings are currently supported by Das and Desh, Mapson and Gimnez [3, 6, 12] suggesting a decrease in ascorbic acid may be due to oxidation of ascorbic acid in dehydro ascorbic acid ascorbic acid with the enzyme ascorbinase. Similar results were also reported by Tarkase and Desai [17] in mosambi.




Fig 8: Effect of different treatment on the ascorbic acid of fruit

Conclusion

It is concluded that among all treatments, T₄ (HDPE perforated) gave better results in traits like phenol content, TSS, total terrible acid, total sugars, reducing sugars and non-reducing sugars, as compared to other treatments studied. On the other hand, fruit pH and ascorbic acid was observed better in T₆ and T₃ respectively. As a result of the above discussion, it was suggested that the use of HDPE perforated packaging material might be effective and economical treatment for improving the quality of Mosambi fruit.

References

1. Ali MA, Raza H, Khan MA, Hussain M. Effect of Different Periods of Ambient Storage on Chemical Composition of Apple Fruit. *International Journal of Agriculture and Biology*. 2004;6(2):568-571.
2. Curk F, Ollitrault F, Garcia-Lor A, Luro F, Navarro L, Ollitrault P. Phylogenetic origin of limes and lemons revealed by cytoplasmic and nuclear markers. *Annals of botany*. 2016;117(4):565-583.
3. Das RC, Desh JC. The Effect of Was Elusion and Growth Regulators on the Storage Behavior of Mosambi Fruits. *International Symposium on Sub-trop Hort*; c1967. p. 104-107.
4. D'Aquino S, Molinu MG, Piga A, Agabbio M. Influence of film wrapping on quality maintenance of Salustiana orange under shelf-life conditions. *Italian Journal of Food Science*. 2001;1(13):87-100.
5. Echeverria E, Valich J. Enzymes of Sugars and Acid Metabolism In Stored Valencia Oranges. *Journal of the American Society for Horticultural Science*. 1989;114(3):445-449.
6. Gimnez M, Olarte C, Sanz S, Lomas C, Echavarri L, Ayala F. Influence of Packaging Films on The Sensory And Microbiological Evolution of Minimally Processed Borage (*Borago officinalis*). *J Food Sci*. 2003;68:1051-1108.
7. *Horticultural Statistics At A Glance*; c2018.
8. Ingel HV, Vivas L, Bertuzzi MA, Apella MC, Audisio MC. Edible Films with Anti-Listeria Monocytogenes Activity. *International Journal of Food Science & Technology*. 2010;45(7):1443-1449.
9. Jawanda JS, Singh R, Vij VK. Studies on Extending Post Harvest Life of Kinnow Mandarin. *Punjab Horticultural Journal*. 1978;18(3&4):149-153.
10. Kaur S, Singh S. Effect of Various Post-Harvest Treatments on Shelf Life of Lemon (*Citrus limon*) Cv. Baramasi during Ambient Storage. *Progressive Agriculture*. 2012;12(2):277.
11. Keditsu R, Sema A, Maiti CS. Effect of Modified Packaging and Low Temperature on Post-Harvest Life of Khasimandarin. *J Food Sci. Tech*. 2003;40:646-651.
12. Mapson LW. Vitamins in Fruits. In *Biochemistry and fruits and their products*. Hulme, A.C. (ed). Academic Press, London and New York; c1970. p. 1.
13. Peterson JJ, Beecher GR, Bhagwat SA, Dwyer JT, Gebhardt SE, Haytowitz Rab A, *et al*. Fruit Quality and Senescence Related Changes In Sweet Orange Cultivar Blood Red Uni-Packed In Different Packing Materials. *Sarhad Journal of Agriculture*. 2010;26(2):221-228.
14. Saltveit ME. Is It Possible To Find an Optimal Controlled Atmosphere. *Postharvest Biology and Technology*. 2003;27:3-13.
15. Mohd S, Khan A, Kumar M. Effect of Storage on Some Biochemical Parameters of Selected Fresh Fruits Juice. *International Journal of Pharma and Bio Sciences*. 2013;4(2B):659-663.
16. Tarkase BG, Desai UT. Effect of Packaging and Chemicals on Storage of Orange Cv. Mosambi. *Journal of Maharashtra Agricultural Universities*. 1989;14(1):10-13.
17. Vines HM, Oberbacher MF. Changes in Carbon Dioxide Concentrations within Fruit and Containers during Storage. *Proceedings of the Florida State Horticultural Society*. 1961;74:243-246.