



E-ISSN: 2663-1067  
P-ISSN: 2663-1075  
<https://www.hortijournal.com>  
IJHFS 2023; 5(1): 101-105  
Received: 05-04-2023  
Accepted: 07-05-2023

**Shreejan Kumar Pandey**  
Mahendra Ratna Multiple  
Campus Ilam, IAAS,  
Tribhuvan University,  
Kathmandu, Nepal

**Prashna Budhathoki**  
Mahendra Ratna Multiple  
Campus Ilam, IAAS,  
Tribhuvan University,  
Kathmandu, Nepal

**Sanu Krishna Maharjan**  
Department of Horticulture,  
Institute of Agriculture and  
Animal Science, Tribhuvan  
University, Kathmandu,  
Nepal

**Corresponding Author:**  
**Shreejan Kumar Pandey**  
Mahendra Ratna Multiple  
Campus Ilam, IAAS,  
Tribhuvan University,  
Kathmandu, Nepal

## Soil status and performance of Nepal cardamom (*Amomum subulatum* Roxb.) under different land uses

**Shreejan Kumar Pandey, Prashna Budhathoki and Sanu Krishna Maharjan**

DOI: <https://doi.org/10.33545/26631067.2023.v5.i1b.168>

### Abstract

Soil fertility evaluation is the most fundamental decision-making tool for effective sustainable soil nutrient management of a particular area. Soil properties vary between land use systems and understanding the variation in soil properties within land use systems is crucial. Thus, the study was conducted from March 2022 to April 2022 in Phaktalung, Taplejung of Nepal for evaluation of the soil nutrient conditions of different land use systems in Randomized Complete Block Design. Different land use systems i.e., Nepal Cardamom orchard in khet land, Nepal Cardamom orchard in bari land, Nepal Cardamom orchard in Pakho land were taken as treatments. A total of 27 samples from cardamom orchard of Ramsai variety were randomly collected, 9 samples each from khet land, bari land and Pakho land respectively at a depth of 0-30 cm and are analyzed. Research analysis revealed that the soil organic matter, nitrogen, potassium was found maximum in Treatment 3 (Pakho land) which is 4.12%, 0.20% and 331.5 kg/ha respectively. Phosphorous (51.10 kg/ha) was found maximum in Treatment 2 which was Bari land. Whereas cardamom yield (0.55 Mt/ha), number of tillers (29.77) were found maximum in Treatment 1 which was khet land. While soil pH, diameter of tillers, length of tillers, number of inflorescences was found non-significant. Soil texture was found to be sandy loam type.

**Keywords:** Cardamom, soil, land, khet, Bari, pakho

### 1. Introduction

Nepal cardamom (*Amomum subulatum* Roxb.) is one of the world's ancient species belongs to family Zingerberaceae, popularly known as Alainchi in Nepali. It is also known as "Queen of spices". It mainly grows wild in forest ecosystem and is also domesticated in sub-himalayan region, at altitude ranging from 1000 to 2200 m above mean sea level. It is perennial herb and shade loving plant which requires well distributed rainfall of about 3000-5000 mm/year and generally grown in tracts (Gudade, *et al.*, 2013; Gupta, 1983; Singh D., 1978; Subba, 1984) [6, 7, 20, 22]. Nepal cardamom is also known as black cardamom, large cardamom, big cardamom (Shrestha, *et al.*, 2018) [18]. This plant is 1.5 to 3.0 m tall that consists of subterranean rhizomes and several leafy aerial shoots/tillers (Bisht, Negi, Bhandari, & Sundriyal, 2011) [3]. It is commonly cross-pollinated crop, although self-fertilization can also occur. The plant has several tillers consisting of pseudo stems with leaves on upper part and inflorescence (spike) appears on the rhizome from the point where pseudo stem shoots up and traditional kilns are used to cure harvested capsules (Sharma E., Sharma, Singh, & Sharma, 2000) [16].

In recent years, Nepal cardamom has been a major agricultural cash crop and export commodity of Southeast Asian countries (Sharma, Sharma, & Sharma, 2009) [17]. It is mostly grown in the eastern hills of Nepal namely Panchthar, Taplejung, Tehrathum, Sankhuwasabha, Bhojpur (Pun, 2019). The dried fruits of Nepal cardamom is widely used in foods, beverages, perfumes, and medicines which is a high-value, low-volume spice crop (Singh & Pothula, 2013) [19].

Nepal cardamom can be grown in the sandy loam soil, slightly acidic in nature with gentle sloping land of about 5 - 8° provide highly suitable conditions (Baniya, Böehme, & Baniya, 2009) [2]. According to Dutta & Dutta (1997) [4], soil health is an important parameter to promote the healthy and vigorous growth of the plants.

Soil properties are greatly impacted by different land use and management practices and for determining production constraints related to soil nutrients, it is essential to have knowledge of variation in soil properties within farmland use (Spurgeon, Keith, Schmidt, Lammertsma, & Faber, 2013) [21]. It is estimated that 60% soil of Nepal have low organic matter (OM), 23% have low phosphorous (P), 18% have low potassium (K), and 67% of the soils are acidic (Mandal, 2016) [11].

Soil fertility is the sum of physical, chemical, and biological factors that affect soil structure, porosity, availability of macro and micro nutrients, and the activity of biological organisms which influence crop growth and impose limitations on crop yield (Rowell, 2014) [15]. Soil fertility plays important role for making soil alive, thus for sustainable soil management and crop production. Evaluation of soil fertility is now becoming a routine work. Among various techniques for soil fertility evaluation, soil testing is an indispensable tool in soil fertility management for sustained soil productivity (Havlin, Tisdale, Nelson, & Beaton, 2010) [1].

## 2. Materials and Methods

**2.1 Site description:** The experiment was carried out in Phaktalung Rural Municipality of Province 1 representing high hills of Nepal from March 2022 to April 2022 the important area under Cardamom Zone of Taplejung district. Khet lands are lowland levelled terraces. Bari lands are upland sloping terraces and Pakho land are forest land which are sloppier than bari and normal cultivation of major crops is not possible due to its topography.

### 2.2 Treatment details

The experiment was conducted in Randomized Complete Block Design (RCBD) with 3 treatments namely Treatment 1: Orchard in khet land, Treatment 2: Orchard in bari land, Treatment 3: Orchard in Pakho land were taken as treatments. From each treatment, 9 composite samples of soil (as replications) were obtained. Nine Sampling plots (each with 400 square m.) at random locations for each treatment were selected from selected locations. Altogether, there were 27 sampling plots (1 treatment=9 sampling plots).

**2.3 Soil Sample collection:** Soil samples were collected from each treatment following Simple Random Sampling Technique. Firstly, plots were subdivided and then samples were taken through lottery method of Simple Random Sampling. Soil at the depth of 30 cm below the surface was collected by making 'V' shape cut. The starting point was selected randomly. Samples were taken from each land uses with Zigzag method. Soil samples were collected from nine replicated sites of each land uses type within the study area. Altogether 27 samples were collected randomly for analysis of different soil parameters. Various instruments were used in order to collect soil samples in the field and to measure the physio-chemical properties of soil in the laboratory. Spades were used in accordance with the type of land for collection of soil samples. Similarly, soil-packing polythene was used to place the soil samples.

**2.4 Laboratory analysis:** Different chemicals and lab equipment were used in lab during qualitative as well as quantitative data evaluation of soil. The collected soil

samples were analysed at Regional Soil Laboratory, Jhumka, Sunsari. All soil physio-chemical data including pH, Soil Organic Matter, Nitrogen, Phosphorus and Potassium were quantitative data whereas soil texture was qualitative. The different soil parameters that were tested as well as methods that were adopted during laboratory analysis are shown on Table 1.

**Table 1:** Parameters and methods adopted for the laboratory analysis

Parameters	Analysis methods
Soil organic matter	Walkey and Black Method (Walkey & Black, 1934) [32]
Total nitrogen in soils	Walkey and Black Method (Walkey & Black, 1934) [32]
Available Phosphorus	Modified Olsen's Bicarbonate Method (Olsen <i>et al.</i> , 1954) [25]
Available Potassium	Flame Photometer Method (Barnes <i>et al.</i> , 1945) [26]
Soil texture	Hydrometer Method (Gee & Bauder, 1986) [27]
Soil pH	Electrometric method

### 2.5 Soil rating

Soil organic matter (SOM%), total N (%), available P (%) and available K (%) were rated according to the rating chart provided by Regional soil laboratory, Jhumka, Sunsari (Table 2) and the soil pH was rated by the rating chart provided by Khatri-Chhetri (1991) [33] (Table 3)

**Table 2:** Rating chart for classification of fertility status of the studied soils according to Regional Soil Laboratory, Jhumka, Sunsari

Nutrient status	SOM (%)	Total N (%)	Available P (mgkg <sup>-1</sup> )	Available K (mgkg <sup>-1</sup> )
Low	<= 2.5	<= 0.10	<= 30	<= 110
Medium	2.51 - 5.0	0.11-0.20	30.1-55	110.1 - 280
High	> 5.0	> 0.20	> 55	> 280

**Table 3:** Rating chart for soil reaction according to Khatri- Chhetri (1991) [33]

Soil pH value	Soil reaction rating
<6	Acidic
6.0-7.5	Neutral
>7.5	Alkaline

### 2.6 Parameters recorded

#### 2.6.1 Data collection and analysis

**2.6.1.2 Number of tillers:** Three cardamom bushes were selected randomly and number of tillers present in each bush were counted during soil sample collection time.

**2.6.1.3 Number of inflorescences:** One cardamom bush was selected randomly. Then five tillers from selected bush were selected randomly and number of inflorescences present in each tiller were counted during soil sample collection time.

**2.6.1.4 Pseudo stem height:** Five cardamom bushes were selected randomly. One tillers from each bush was selected randomly then pseudo stem height of each tiller was measured with the help of measuring tape during soil sample collection time.

**2.6.1.5 Pseudo stem diameter:** Five cardamom bush were

selected randomly. One tiller from each bush was selected randomly, then pseudo stem diameter of each tiller was measured with the help of digital Vernier calliper during soil sample collection time. Measurement of pseudo stem height and diameter were taken from same plant.

The information on Production was obtained from the survey and was correlated with the result of soil parameters obtained from lab analysis.

The data were analyzed using Duncan's Multiple Range Test (DMRT) at 5% level of significance for the separation of means using the package agricolae. The collected data were entered into MS-excel version 2019 and were analyzed using R studio version 3.5.1.

### 3. Results and Discussion

#### 3.1 Soil Organic Matter

Analysis of variance revealed that the soil organic matter of cardamom field was significantly different (Table 4). Highest amount of soil OM was found in Pakho land (4.123%) and lowest amount of soil organic matter was found in khet land (2.981%) where the soil organic matter of Bari land (3.500%) was between above two. This result agrees with that of Ghimire, Bhatta, Pokhrel, Kafle, & Paudel (2018) [28] who reported that the forest soil had the highest SOC, followed by Bari and khet. Continuous cultivation with minimum addition of SOM in sandy textured soil cause lower SOC content in Khet and Bari soil (Paudel & Thapa, 2001) [12]. The results were also obtained similar to Kharal, Khanal, & Panday (2018) [29] who reported that highest amount of soil organic matter was found in Pakho land followed by Bari and khet land. Tillage improves the aeration of soil which speeds up decomposition of OM as results of microbial action and often increases erosion (Funderburg, 2016) [5]. Hence, the Pakho land showed higher soil OM due to high OM accumulation and no-tillage disturbance as compared to bari

and khet land.

#### 3.2 Soil Nitrogen

The results of the study indicated that the effect of different land uses on soil nitrogen was highly significant. The total nitrogen was highest in Pakho land (0.2067%) whereas lowest N percentage was reported in khet land (0.150%), whereas total nitrogen percentage in bari land (0.175%) was reported in between two. The results were obtained similar to Kharal, Khanal, & Panday (2018) [29] who reported that the highest soil N was observed from forest land and the lowest was observed in the lowland (khet) and upland (*bari*) consist nitrogen in-between. There was a positive correlation ( $r = 0.998$ ) between SOM and N level indicating that the highest nitrogen level was in Pakho land. Similar findings were put forward by Charan, *et al.* (2013) [30]. They found that amount of nitrogen increases with increases in SOM because soil required nitrogen to decompose the organisms into organic matter. Low soil OM and tillage might be the reason for lower N contents in bari land and khet land as soil tillering increases the susceptibility to erosion (Funderburg, 2016) [5].

#### 3.3 Available Phosphorous

The results of the study indicated that the effect of different land uses on available phosphorous was highly significant. The available phosphorous was found highest in bari land (51.10 kg/ha) followed by khet land (42.07 kg/ha). The lowest available phosphorous was reported in Pakho land (36.30 kg/ha). The results were obtained similar to Kharal, Khanal, & Panday (2018) [29] who reported that the available phosphorous was found highest in upland (*bari*) and lowest in Pakho land where lowland (khet) consist available phosphorous in-between. In cultivated soil availability of phosphorous strongly depends on soil pH (Lindsay, 1979) [10]. Hence, low P in Pakho land might be due to low pH.

**Table 4:** Soil organic matter, Nitrogen, Phosphorous, Potassium and soil pH under different land uses of cardamom orchards in Taplejung, Nepal

Treatments	Organic matter (%)	Nitrogen (%)	Phosphorous (kg/ha)	Potassium (kg/ha)	pH
Khet	2.981 <sup>c</sup>	0.1500 <sup>c</sup>	42.07 <sup>b</sup>	191.7 <sup>c</sup>	5.244
<i>Bari</i>	3.500 <sup>b</sup>	0.1756 <sup>b</sup>	51.10 <sup>a</sup>	282.9 <sup>b</sup>	5.378
Pakho	4.123 <sup>a</sup>	0.2067 <sup>a</sup>	36.30 <sup>c</sup>	331.5 <sup>a</sup>	5.211
LSD (=0.05)	0.4382	0.021	4.547	41.28	0.1978
CV (%)	12.4	11.7	10.5	15.4	3.7
Significance	***	***	***	***	ns
S.Em (±)	0.1462	0.007	1.517	13.77	0.0660
Grand mean	3.535	0.18	43.16	268.7	5.278

**Note:** Means followed by the same letter (s) or without any letter in a column are not significantly different at 5% level of significance in DMRT test. (S.Em – Standard Error of mean, CV – Coefficient of Variation, LSD – Least Significance Difference, NS-Non-significant, \*\*\* is significant at  $p < 0.001$ ).

#### 3.4 Available Potassium

The results of the study indicated that the effect of different land uses on available potassium was highly significant. The available potassium was found highest in Pakho land (331.5 kg/ha) followed by bari land (282.9 kg/ha). The lowest available potassium was reported in khet land (191.7 kg/ha). The results were obtained similar to Kharal, Khanal, & Panday (2018) [29] who reported that the forest land had higher K level and lowland (khet) had lowest K level where upland (*bari*) consist potassium level in-between. Due to higher leaching loss and more K harvest from soils there is low level of K in low land (khet). Under irrigated

conditions, K is subjected to considerable leaching loss (Harter, 2007) [8].

#### 3.5 Soil pH

The results of the study indicated that there was no significant effect of different land uses on soil pH. The soil pH was found highest in bari land (5.378) followed by khet land (5.244). The lowest soil pH was found in Pakho land (5.211). The results were obtained similar to Kharal, Khanal, & Panday (2018) [29] who reported that the Soil pH was found lowest in forest land and highest in upland (*bari*) where pH of lowland (khet) lies in-between. According to

the rating chart adopted from (khatri-chhetri, 1991) [33], soil reaction was found acidic in all three land uses in cardamom field of taplejung. The lower soil pH in forest (Pakho) land might due to its higher slope (Yeshaneh, 2015) [23] and might also due to high OM content (Harter, 2007) [8]. Low pH in all land uses may be due to higher rainfall in eastern

part of Nepal.

### 3.6 Soil Texture

The soil texture observed during the soil physical property analysis was found to be sandy loam type (Table 5).

**Table 5:** Soil texture with the proportions of sand, silt and clay.

Treatments	Clay (%)	Sand (%)	Silt (%)	Textural class
Khet	2.244444444	60.64444444	37.11111111	SL
Bari	2.911111111	63.08888889	34	SL
Pakho	2.022222222	72.2	25.77777778	LS

**Note:** SL represents sandy loam type, LS represents loamy sand type

### 3.7 Cardamom Performance

#### 3.7.1 Yield

The results of the study indicated that the effect of different land uses on Cardamom performance (Yield) was highly significant. The yield was found highest in khet land (0.5578 Mt per ha) followed by bari land (0.3756 Mt per ha). The yield was found lowest in Pakho land (0.2144 Mt per ha). Although the nutrient was found higher in Pakho land yield was found higher in khet land. Highest yield of cardamom on khet land might be due to higher rates of nutrient cycling on low land areas, where organic matter decomposes more rapidly, releasing essential nutrients for plant uptake & also, it can be due to more direct sunlight compared to upland and forested regions, which can promote higher rates of photosynthesis and biomass production. However, other factors such as irrigation facility, soil moisture resin, depth of soil, root zone and other nutrients etc. also need to be considered in the nutrient balance. According to Rollon (2021) [31] high number tillers produce more yields and higher number of tillers were found in khet land.

#### 3.7.2 Number of Tillers

The results of the study indicated that the effect of different land uses on Cardamom performance (Number of tillers) was highly significant. The Number of tillers were found highest in khet land (29.77) followed by bari land (24.91). The number of tillers were found lowest in Pakho land

(22.28). In cardamom tillage is not common as it is perennial crop. Higher effective tillers in no tillage land might be due to favourable growing conditions and higher carbon accumulation in soil (Rahman, Biswas, Maniruzzaman, Choudhury, & Ahmed, 2017) [14].

#### 3.7.3 Diameter

The results of the study indicated that there was no significant effect of different land uses on diameter of tillers. The diameter was found highest in bari land (19.58mm) followed by Pakho land (19.44 mm). The diameter of tillers was found lowest in khet land (19.10 mm).

#### 3.7.4 Length of Tillers

The results of the study indicated that there was no significant effect of different land uses on length of tillers. The length of tillers was found highest in khet land (213.0) followed by bari land (209.6). The length of tillers was found lowest in Pakho land (207.0).

#### 3.7.5 Number of Inflorescences

The results of the study indicated that there was no significant effect of different land uses on number of inflorescences. The number of inflorescences was found highest in bari land (3.022) followed by Pakho land (2.978). The number of inflorescences was found lowest in khet land (2.967).

**Table 6:** Yield, Number of tillers, Diameter, Lengths of tillers and number of inflorescences under different land uses of cardamom orchards in Taplejung, Nepal

Treatments	Yield (mt/ha)	Number of tillers	Diameter (mm)	Lengths of tillers(cm)	Number of inflorescences
Khet	0.5578 <sup>a</sup>	29.77 <sup>a</sup>	19.10	213.0	2.967
Bari	0.3756 <sup>b</sup>	24.91 <sup>b</sup>	19.58	209.6	3.022
Pakho	0.2144 <sup>c</sup>	22.28 <sup>c</sup>	19.44	207.0	2.978
LSD (=0.05)	0.0842	0.855	2.211	14.74	0.5041
CV (%)	22	3.3	11.4	7.0	16.9
Significance	***	***	ns	ns	ns
S.Em (±)	0.0281	0.285	0.737	4.92	0.1681
Grand mean	0.383	25.65	19.37	209.9	2.989

**Note:** Means followed by the same letter (s) or without any letter in a column are not significantly different at 5% level of significance in DMRT test. (S.Em – Standard Error of mean, CV – Coefficient of Variation, LSD – Least Significance Difference, NS-Non-significant, \*\*\* is significant at  $p < 0.001$ ).

### 4. Conclusion

The result of present findings concluded that the different land uses of cardamom orchard had a considerable effect on soil nutrient pool except for soil pH and also had effect on cardamom performance i.e., Yield and Number of tillers. Soil organic matter, nitrogen, and potassium were found higher in Pakho land, phosphorous was found higher in Bari

land. Yield, Number of tillers and length of tillers were found higher in khet land, diameter and number of inflorescences were found higher in Bari land. In the study area, significant association was found between the soil organic matter and nitrogen content. From the study, it can be concluded that for enhancing efficacy of cardamom production, timely test of the soil nutrient status of the

cardamom orchards and application of the fertilizers based on the soil test results should be conducted by farmers rather than based on blanket recommendations. Nutrients status of other micronutrients should also be assessed for the overall improvement of cardamom orchards.

## 5. References

- Havlin JL, Tisdale S, Nelson W, Beaton J. Soil Fertility and Fertilizers: An Introduction to Nutrient management (7th Edition ed.). New Delhi: PHI Learning Private Limited; c2010.
- Baniya N, Böehme MH, Baniya S. Physical land suitability assessment for the large cardamom *Amomum subulatum* Roxb. Cultivation in hills of Kathmandu Valley. Chinese Journal of Population Resources and Environment. 2009;7(4):59-63.
- Bisht V, Negi J, Bhandari A, Sundriyal R. *Amomum subulatum* Roxb: Traditional, phytochemical and biological activities-An overview. African Journal of Agricultural Research. 2011;6(24):5386-5390.
- Dutta AC, Dutta T. Botany for degree students (6th Edition ed.). Bombay, Calcutta and Madras: Oxford University Press; c1997.
- Funderburg E. Organic matter serves important role in soil health. Noble Research Institute: Ardmore, OK, USA; c2016.
- Gudade B, Chhetri P, Deka T, Gupta U, Vijayan A, *et al.* Large cardamom (*Amomum subulatum* Roxb): A spice crop with multiple uses. Indian Journal of Arecanut, Spices and Medicinal Plants. 2013;15(4):15-17.
- Gupta P. Export potential in large cardamom. Cardamom; c1983.
- Harter RD. Acid soils of the tropics. ECHO Technical Note, ECHO; c2007, 11. Retrieved from [http://courses.umass.edu/psoil370/Syllabus-files/Acid\\_Soils\\_of\\_the\\_Tropics.pdf](http://courses.umass.edu/psoil370/Syllabus-files/Acid_Soils_of_the_Tropics.pdf)
- Dhakal SR, Mandal RA, Mathema AB. Soil nutrients and carbon increment dynamics in broadleaf and chirpine forests of community forests Bhaktapur, Nepal. International Journal of Geography, Geology and Environment. 2021;3(2):35-46.
- Lindsay WL. Chemical equilibria in soils. John Wiley and Sons Ltd; c1979.
- Mandal S. National workshop on sustainable soil management program. Lalitpur: Soil Management Directorate; c2016.
- Paudel GS, Thapa GB. Changing farmers' land management practices in the hills of Nepal. Environmental Management. 2001;28:789-803.
- Pun AB. A Review on Different Factors of Large Cardamom Decline in Nepal. Asian Journal of Research in Crop Science; c2019. p. 1-6. DOI:10.9734/AJRCS/2018/46732
- Rahman M, Biswas J, Maniruzzaman M, Choudhury A, Ahmed F. Effect of tillage practices and rice straw management on soil environment and carbon dioxide emission. The Agriculturists. 2017;15(1):127-142.
- Rowell DL. Soil science: Methods & applications. New York; c2014. Routledge. DOI: <https://doi.org/10.4324/9781315844855>
- Sharma E, Sharma R, Singh K, Sharma G. Large Cardamom Farming in the Sikkim Himalaya. Mountain Research and Development. 2000;20(2):108-111.
- Sharma G, Sharma R, Sharma E. Traditional knowledge systems in large cardamom farming: biophysical and management diversity in Indian mountainous regions. Indian Journal of Traditional Knowledge (IJTK). 2009;8(1):17-22.
- Shrestha J, Prasai H, Timsina K, Shrestha K, Pokhrel D, Poudel K, *et al.* Large cardamom in Nepal: Production practice and economics, processing and marketing. Pakhribas, Dhankuta: National Commercial Agriculture Research Program, Nepal; c2018.
- Singh AI, Pothula AK. Postharvest processing of large cardamom in the eastern Himalaya. Mountain Research and Development. 2013;33(4):453-462.
- Singh D. Large cardamom. Cardamom; c1978. p. 3-13.
- Spurgeon DJ, Keith AM, Schmidt O, Lammertsma DR, Faber JH. Land-use and land-management change: relationships with earthworm and fungi communities and soil structural properties. BMC Ecology. 2013;13:1-13.
- Subba J. Agriculture in the hills of Sikkim, Sikkim Science Society. Gangtok; c1984.
- Yeshaneh GT. Effect of slope position on soil physico-chemical properties with different management practices in small holder cultivated farms of Abuhoy Gara Catchment, Gidan District, North Wollo. African Journal of Environment Protection. 2015;3(5):174-179.
- Chandravanshi MD, Patel JR, Kasyap S. Effect of integrated nutrient management on growth and yield of Urdbean (*Vigna mungo* L.) in Chhattisgarh plain. International Journal of Advanced Chemistry Research. 2022;4(2):260-265. DOI: 10.33545/26646781.2022.v4.i2d.109
- Olsen SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. US Department of Agriculture; c1954.
- Barnes RB, Richardson D, Berry JW, Hood RL. Flame photometry a rapid analytical procedure. Industrial & Engineering Chemistry Analytical Edition. 1945;17(10):605-11.
- Gee GW, Bauder JW. Particle-size analysis. Methods of soil analysis: Part 1 Physical and mineralogical methods. 1986;5:383-411.
- Ghimire P, Bhatta B, Pokhrel B, Kafle G, Paudel P. Soil organic carbon stocks under different land uses in Chure region of Makawanpur district, Nepal. SAARC Journal of Agriculture. 2018;16(2):13-23.
- Kharal S, Khanal BR, Panday D. Assessment of soil fertility under different land-use systems in Dhading District of Nepal. Soil Systems. 2018;2(4):57.
- Charan G, Bharti VK, Jadhav SE, Kumar S, Acharya S, Kumar P, *et al.* Altitudinal variations in soil physico-chemical properties at cold desert high altitude. Journal of Soil Science and Plant Nutrition. 2013;13(2):267-77.
- Suwa R, Rollon R, Sharma S, Yoshikai M, Albano GM, Ono K, *et al.* Mangrove biomass estimation using canopy height and wood density in the South East and East Asian regions. Estuarine, Coastal and Shelf Science. 2021;248:106937.
- Griffiths PH, Walkey WA, Watson HB. 131. The nitration of anisole. Journal of the Chemical Society (Resumed); c1934. p. 631-633.
- Khatri-Chhetri T. Introduction to soils and soil fertility. Tribhuvan University Institute of Agricultural and Animal Science, Rampur, Chitwan, Nepal; c1991. p. 233.