



E-ISSN: 2663-1067
 P-ISSN: 2663-1075
 IJHFS 2020; 2(2): 25-35
 Received: 18-06-2020
 Accepted: 24-07-2020

Muhammad Arshad Ullah
 Pakistan Agricultural
 Research Council, Islamabad,
 Pakistan

Boosting up the economy of Pakistan through agarwood production

Muhammad Arshad Ullah

Abstract

Aquilaria species have been the primary source for agarwood and are distributed eastward from India (Bengal and Assam) to the island of New Guinea and southward from tropical China. Agarwood is dark resinous heartwood that forms in *Aquilaria* and *Gyrinops* trees in the plant family Thymelaeaceae. Agarwood has been used for medicinal purposes for thousands of years, and continues to be used in Ayurvedic, Tibetan and traditional East Asian medicine. High-grade agarwood powder is used for aromatherapy and also used in the production of pharmaceutical tinctures. The number and types of agarwood metabolite constituents of each reported studies vary depending on the agarwood source, extraction methods and analysis approaches used. Agarwood, also known as eaglewood or gaharu, is a valuable non-timber forest product which sometimes grows in *Aquilaria* species. The genus species occur mainly in South and Southeast Asia. As a result of a defense mechanism to fend off pathogens, *Aquilaria* species develop agarwood which can be used for incense, perfume, and traditional medicines. The main markets for these products are in South and East Asia and the Middle East. The high prices demanded for agarwood has led to the rapid depletion of *Aquilaria* trees in natural forests. The search for agarwood has spread from one country to another. Efforts have been undertaken to increase the production of the infected wood by deliberately wounding the trees. A variety of methods is used towards this end. Some recently developed techniques have proven to be most effective. This resulted in planting of *Aquilaria* trees by small holders as well as large industrial size plantations. In this paper, we shall discuss all the concerns of agarwood for better yield, quality and maximum utilities for humanity as well as the environment safety to enhance the socio economic condition of the Pakistan.

Keywords: domestication, future developments, plantations, pakistan, socio-economic

Introduction

Aquilaria species have been the primary source for agarwood and are distributed eastward from India (Bengal and Assam) to the island of New Guinea and southward from tropical China. The resin, is widely used for the production of incense and perfume (essential oils) and for medical purposes (Mohamed and Lee, 2016; Rhind, 2014) ^[65, 91]. The valuable resin develops and accumulates in the stem and branches of the tree after a process of injury and fungal infection (Donovan and Puri, 2004; Rasool and Mohamed, 2016; Sangareswari *et al.*, 2016) ^[36, 90, 95], but this process occurs in only a few trees in wild populations (Paoli *et al.*, 2001) ^[81]. The demand for and trade in agarwood continues today, and with increasing wealth in consumer countries over recent decades, demand exceeds supply (Barden *et al.*, 2000; Compton and Ishihara, 2006) ^[15, 32]. This has resulted in increased prices (Wyn and Anak 2010) ^[116], natural resource decline (Zhang *et al.* 2008) ^[122], reduction in product quality (Antonopoulou *et al.*, 2010) ^[8], increasing interest in cultivation (Hoang and Nghi, 2011; Page and Awarau, 2012; Persoon, 2008; Rahman *et al.*, 2015) ^[49, 80, 84, 88], and developing methods for resin induction (Liu *et al.*, 2013; Mohamed *et al.*, 2014; Persoon, 2008) ^[67, 84, 77]. In Japan, it is known asjinkoh (also means the sinking incense) (Brechtbill, 2012) ^[22]. In European languages such as Portuguese, agarwood is known asaguilaorpaod' aguila; in French, the wood is recognized as d'aigle; and in English as eaglewood. The latter is taken from Medieval Latin (*aquilaria*) which is derived from the Greek (agallochum). The Sanskrit word for incense is known as agar bhatti, which is derived from the word for agarwood (agāru) (McKenna and Hughes, 2014) ^[72].

Agarwood Manufacturing

Agarwood is dark resinous heartwood that forms in *Aquilaria* and *Gyrinops* trees in the plant family Thymelaeaceae (Blanchette, 2006) ^[18].

Corresponding Author:
Muhammad Arshad Ullah
 Pakistan Agricultural
 Research Council, Islamabad,
 Pakistan

It is highly prized that is extremely rare. It has at least a 3000 years history in the Middle East, India, China and Japan (Le, 2003) ^[64]. In Japan, agar incense is more expensive than gold (Japan Talk, August 26, 2012). The value of agarwood shipped out of Singapore alone each year has been estimated to exceed \$ 1.2 billion (Hansen, 2000) ^[46]. This aromatic resinous wood has many common names including agarwood, gaharu, eaglewood, aloeswood, agila wood, aguru, agar, oud, ude, ud, ood, oode, jinkoh, jinko, Ch'Ing Kuei Hsiang, Ch'En Hsiang, Chan Hsiang, Chi Ku Hsiang, Huang Shu Hsiang, kalambak, grindsanah, etc (Blanchette and Van Beek, 2005) ^[20]. Agarwood is a diseased tissue corresponding production of oleoresin which begins to become odoriferous and this aromatic resinous wood is called agarwood (Ba, 2010) ^[12]. At this moment, no other substances are similar as agarwood. Chemical substitutes are already available for perfume but do not come even close in emulating the natural product (Ali, 2006) ^[7].

The identification of the small proportion of the trees having agar is difficult and destructive, which added greatly to the near-extinction of natural stands of trees (Blanchette, 2006) ^[18]. Also, large-scale logging operations have destroyed many forested areas where the *Aquilaria* trees are found. Thus, the current source of agarwood, the naturally-growing old-growth *Aquilaria* trees is becoming extinct (Hansen, 2000) ^[46].

Aquilaria malaccensis is a large, evergreen tree, up to 20 - 40 m tall (Adelina, 2004) ^[2]. This tree has an unusual anatomy and specialized cells within the xylem that produce the resin (Norsuzieana, 2009) ^[79]. Flowers are hermaphroditic, up to 5 mm long, fragrant and yellowish green or white which blooms at June. Fruits which appear in August are green, egg-shaped capsule, leathery covered with fine hairs, 4 cm long, and 2.5 cm wide. There are two seeds per fruit (Sumarna, 2008) ^[105].

Aquilaria is unique in producing phloem bundles within the xylem. This network of phloem and parenchyma produce and distribute the resin around affected areas as a tree defense reaction (Blanchette and Van, 2005) ^[20].

Uses of agarwood

Medicine: Agarwood has been used for medicinal purposes for thousands of years, and continues to be used in Ayurvedic, Tibetan and traditional East Asian medicine. High-grade agarwood powder is used for aromatherapy and also used in the production of pharmaceutical tinctures. It is prescribed in traditional East Asian medicine to promote the flow of qi, relieve pain, arrest vomiting by warming the stomach, and to relieve asthma (Hajar, 2013) ^[45].

Although, several commercial synthetic agarwood fragrance compounds are available, they can produce only low-quality fragrances, owing to the chemical structure of natural agar oil. Agarwood incenses have also been used as a fragrance in soaps and shampoos (Schippmann, 2001) ^[96].

Incense: Agarwood incense is burned to produce a pleasant aroma, its use ranging from a general perfume to an element of important religious occasion. It is highly psychoactive and is used for incense (Qi *et al.*, 2005) ^[87]. Taiwanese consumers purchase agarwood for the manufacture of incense sticks, which are used in prayers during many traditional festivals and ceremonies to bring safety and good luck (TRAFFIC East Asia-Taipei, in litt. to TRAFFIC

International 1, 2 May 2000). In Japan, it is considered by many to be sacred and is used to anoint the dead. In Buddhism, it serves as a major ingredient in many incense mixtures, and it is considered to be one of the three integral incenses, together with sandalwood and cloves (Barden *et al.*, 2000) ^[15].

Economic values

The value of first-grade agarwood is extremely high. It is sold in the form of woodchips, wood pieces, powder, dust, oil, incense ingredients and perfume for several thousand US dollars per kilogram (Barden *et al.*, 2000; Gunn *et al.*, 2004; Compton, 2007) ^[15, 31] which varies with geographical location and cultural deposition. Agarwood chips start at \$30 per kilo up to \$9,000 per kilo depending on how much resin is inside the chips (Babatunde, 2015) ^[13]. First-grade agarwood is one of the most expensive natural raw materials in the world.

Agarwood oil fetches similarly high prices (Agarwood "Wood of Gods" International Conference, 2003). When agarwood chips are processed into oil, the agarwood oil was sold at US \$ 30,000 per kg (Nanyang Siang Pau, 15 August 2005). The current global market for Oud oil and other related agarwood products is estimated to be in the range of US\$ 6 to 8 billion (Akter *et al.*, 2013) ^[5] and the major industry buyer of Oud oil, is expected to exceed US \$ 36 billion in 2017. For Integrated cropping - Set up 1,000 ha plantation for agarwood and banana production internal rate of return (IRR) and benefit-cost ratio (B/C ratio) are 54.85% and >1 (3.30) respectively (Mamat *et al.*, 2010) ^[70].

There are about 100 enterprises producing agar wood and agar oil in Bangladesh (Abdin, 2014) ^[1]. Local entrepreneurs are claiming that this is a 100% export oriented sector based on local raw materials and using indigenous technology. They are exporting about US\$ 62.5 to 125 thousand per year.

The agar tree (*Aquilaria malaccensis*) contains dark resinous heartwood called agarwood, eaglewood, oud, gaharu or aloeswood (Barden *et al.*, 2000) ^[15], and offers a primary source of essential oil called oleoresin. This wood has high demand for medicine, incense and perfumes across Asia, Middle East and Europe (Abdin, 2014) ^[1].

In Bangladesh, the production of agar-wood started about 400 years ago in the Suzanagar union under Barlekha Upazila of Moulvibazar District in Sylhet (Abdin, 2014) ^[1]. About 25,000-30,000 workers were engaged in cultivation, collection, processing and marketing of agar and agar-based products in the country (Baksha, 2009) ^[14]. However, little is known about the economic feasibility of agar production in Bangladesh. Reliable information on the financial performance of agar plantations is also lacking (Rahman *et al.*, 2014; Uddin *et al.*, 2018; Akter *et al.*, 2008; Islam *et al.*, 2014; Akter *et al.*, 2013) ^[10, 4, 5].

Economic Benefits

Agarwood is a precious incense substance which is extremely rare. This fragrant wood is only formed in *Aquilaria* and *Gringos* species under family the Thymelaeaceae (Blanchette, 2006) ^[18]. The more resin deposit, the more quality ensure and the precious wood would be formed. It has been using for 3000 years in the China, Japan, India, and especially in Arabian countries (Le, 2003) ^[64] and known by many names depending on place and cultures such as Agar (Bangladesh, India), Aguru

(Sanskrit), Oud (Arabic), Eagle/Agilawood (Europe) (Akter *et al.*, 2013) ^[5].

By 1970s, it was sold up to \$42.5 and then raised straightly \$1250 and \$2500 per kg in 2000 and 2005, respectively (Abdin, 2014) ^[1]. First-grade agarwood is one of the most expensive natural raw materials in the world. Singapore trades agarwood \$1.2 billion alone each year (Hansen, 2000) ^[46]. The trading value of pure agar oil is \$30,000 to 40,000 per kilogram (Abdin, 2014) ^[1]. Based on resin presence, agarwood chips sell from \$30 to \$10,000 per kilogram (Babatunde, 2015) ^[13]. Akter *et al.*, (2013) ^[5] reported that in the year of 2013, the global market for agar oil another agarwood related products was estimated in the range of \$6 to 8 billion and the major agar oil industrial buyer expects to exceed it up to \$36 billion in 2017 (<http://www.ouddh.com/?cid=2065877>).

Agarwood formation is a result of resin synthesis in the xylem tissues of Aquilaria plant. This resin is a chemical substance which consists of aromatic, sesquiterpenes, fatty acids, chromones, phenolic and especially 3-phenyl-2-butanone and alpha-cubebene (Naef, 2011; Hashim *et al.*, 2014) ^[78, 47]. Artificial agarwood or agar oil formation is quite impossible as it is a composition of numerous volatile and semi-volatile compounds. However, the chemical composition of cultivated agarwood is almost similar with naturally inoculated agarwood (Espinoza *et al.*, 2014) ^[37].

Aquilaria spp. trees are indiscriminately and excessively harvested within all ecological zones, and *Aquilaria crassna* is categorised as 'Critically Endangered' by IUCN (Hilton-Taylor, 2000) ^[48]. Aquilaria trees grow well under optimal conditions and annual growth of over 1 m in height per year is common. For further technical information, reference is made to Species Monograph #12 produced by the Lao Tree Seed Project (Jensen, 2002) ^[56].

The agarwood producing tree species are found from India eastwards to Southeast Asian countries (such as Vietnam, Thailand, Malaysia, and Indonesia) to the island of New Guinea (Zich and Compton, 2002) ^[124]. The production of agarwood is uncertain and it is estimated that only 10% of the Aquilaria trees in the forest may contain agarwood (Donovan and Puri, 2004) ^[36].

In the Middle East, agarwood is a symbol of status, wealth and hospitality. An important use of agarwood is the production of incense, the most prized of all incenses. In Middle East, oudh (Arabic word for agarwood) is burnt to release the unique scent and it is a symbol of status, wealth and hospitality (Chang *et al.* 2002) ^[23]. Agarwood is also used as an ingredient in Ayurvedic, Tibetan, Chinese and other traditional Asian medicine. It is also used in the perfume and toiletry products such as soap and shampoo (Chang *et al.*, 2002) ^[23].

Uses of Agarwood Use

	Use	Remark
Production of Incense	Homes, religious ceremonies, rituals and meditation	Precious incense
Oud (Arabic word for wood)	Symbol of status, wealth and hospitality in Middle East	-
Gaharu chips	homes in Middle East	-
Gaharu oil	in perfume and toiletry products such as soap & shampoo	-
Wood	Ayurvedic, Chinese and other Asian traditional Asian medicine	
Wood	sculptures, beads & boxes	Religious purpose

Source: Chang *et al.*, 2002 ^[23] and Barden *et al.*, 2000 ^[15]

Agarwood is a resin filled wood produced in Aquilaria and Gyrinopstrees with a long history of use as incense, perfume and in traditional medicine. Agarwood, has many different names depending on the region of the world where it is found including agar, aloes wood, eagle wood, jinkoh, oud, gaharu, Ch'en Hsiang and Chen-Xiang (Antonopoulou *et al.*, 2010, CITES 2005, Compton and Ishihara, 2004) ^[8, 33, 29].

The loss of this tree from the landscape has been of great concern and several Aquilaria species have been considered endangered according to the International Union for the Conservation of Nature which lists it on their Red List of Threatened Species (IUCN, 2013) ^[53].

Uses for agarwood and its value: Agarwood is used throughout the world but there are two major regions of consumption, the Middle East and Asian markets (Antonopoulou *et al.*, 2010, CITES 2005, Compton and Ishihara, 2004) ^[8, 33, 29].

An essential oil distilled from agarwood has a very long history of use as perfume. Known in the Middle East as oudh (or oud) it can be used as a pure oil or blended with various other substances. This essential oil is mentioned in the bible several times as "aloes" wood (Musselman 2007). There are also descriptions of its use in ancient Islamic religious documents and in reports of its extensive use as perfume as early as the 13th century (Hansen, 2000) ^[46].

Agarwood has also been used in traditional medicines. It has been reported as a component of many traditional Ayurvedic remedies in the Indian subcontinent as well as being used in Tibetan, Chinese, Malayan and Vietnamese medicine (Barden *et al.*, 2000, Kiet 2003 ^[15, 61], Lim and Anack 2010) ^[15]. More modern studies confirm that agarwood has bioactive products that function as effective anti-microbial compounds, it may have anticancer activity, can be used as an antidepressant and used to promote good health in general (CITES, 2005, Dash *et al.*, 2008, Mei *et al.*, 2008) ^[29, 35, 74].

Field trials over many years using methods that mimic the natural production of agarwood have shown great success and are now being used to produce high quality agarwood in young plantation grown trees in many areas of the Indomalaya region (Blanchette and van Beek, 2005) ^[20].

The resin is chemically complex and consists of sesquiterpenes, chromones, fatty acids and phenolic compounds (Naef 2011, Mei *et al.*, 2010) ^[73, 78]. As many as 70 terpenoids have been identified and more continue to be characterized (Espinoza *et al.*, 2014) ^[37].

Agarwood is a dark coloured, fragrant resin embedded in the otherwise whitish wood of Aquilaria species. It is accumulated in the roots and the trunk of the tree as nodules of varying age, shape, size and commercial quality, and is formed as a result of injury and fungal infection (Blanchette, 2007; Jensen, 2009) ^[19, 56]. Agarwood product is very

common in many religious rituals. Wooden pieces and sculptures, beads and bracelets are highly appreciated goods in South and Southeast Asia. However, the major use of agar wood is in the form of extracted essential oil as perfumes, other cosmetic products and in the incense industry. Agar oil is also used in traditional medicines and wine in China and Korea and in ayurvedic medicine in India (Barden *et al.*, 2000; Chen *et al.*, 2011; Gunn *et al.*, 2003; Persoon, 2007) ^[15, 25, 44, 83]. The popularity of Agarwood products in the international market offers the opportunity for a high earning livelihood.

It was evident from the reconnaissance survey and interaction with the villagers that BKS had all the capital (natural, social, human, physical and financial) as household wealth in varying degrees to pursue different livelihood activities (Sherbinin *et al.*, 2008) ^[98].

Agarwood products were seen as an important strategy to earn a high income by virtue of being expensive and therefore, many villagers were attracted to this livelihood activity (Persoon, 2007) ^[83].

Combining plantation or farm activities with wage labour provided a temporary financial relief especially in times when the demand for cash was high for example, paying school fees and expenditure of customary ceremonies (Koczberski *et al.*, 2001) ^[63].

Trees and native vegetation improve the soil beneath them as organic matter, nitrogen and other nutrient content are comparatively higher (Roy *et al.*, 2010) ^[94] in the plantation area. The soil sample data in the present study was not adequate enough to clearly indicate the trend in tree covered land versus annual cropping land as nitrogen, phosphorus and potassium level was similar in both types of land use. Young age plantations are reported to have no impact on phosphorus content of the soil (Zeng *et al.*, 2010) ^[121]. However, it was indicated in the present study that the pH level was less acidic in tree covered land as compared to cassava and maize cultivated land.

Agarwood is a highly prized non-timber forest product which can be used in fragrances, incense, medicines, aromatherapy and religious ceremonies (CITES, 2004; 2005) ^[30]. The value of agarwood shipped out of Singapore alone each year has been estimated to exceed \$1.2 billion (Hansen, 2000) ^[46]. The most important source of agarwood is the Aquilaria spp. tree, which is an angiosperm within the Thymelaeaceae family (Rogers, 2009) ^[93].

In fact, Hong Kong is literally named the "Fragrant Harbour" as it was one of the significant trading centers for agarwood and incense in the early days, when Aquilaria sinensis was widely traded in Hong Kong and then transported to numerous Chinese provinces, Southeast Asia as well as Arabia, along with other incense and spices (Fu, 2008) ^[39].

The healthy wood of Aquilaria trees is white, soft and without scented resins. In a natural environment, agarwood forms only when affected by certain external factors, such as lightning strike, animal grazing, insect attack or microbial invasion, typically around wounded or rotting parts of the trunk (Blanchette and Heuveling, 2009; Pojanagaroon, and Kaewrak, 2006) ^[21, 85].

Agarwood formation occurs slowly and infrequently in old trees and the supply of agarwood from wild sources is far less than market demand. Because of its immense value and rarity, indiscriminate cutting of trees and overharvesting in hope of finding the treasured resin has led to the depletion

of wild trees (Zhang *et al.*, 2008; Soehartono *et al.*, 2000) ^[122, 101]. Eight Aquilaria species, including A. sinensis, were listed on the IUCN red list as endangered species (IUCN, 2008) ^[51].

Gonystylus genus has 20 species, scattering in Southeast Asia regions until Solomon and Nicobar archipelago. *Gyrinops* genus has seven species. Six out of seven are found in East part of Indonesia and one species encountered in Srilanka (Aswandi, 2009) ^[9].

The result showed that their relative population in Sumatera, Kalimantan, Nusa Tenggara, Sulawesi, Maluku and Papua was 26, 27, 5, 4, 6 and 37% respectively (Adijaya, 2009) ^[3]. In order to anticipate the increasing demand and to evade agarwood in nature to become extinct, several efforts to cultivate trees producing agarwood have started to develop in many areas of Indonesia such as North Sumatera, Riau, Jambi, West Java, West Nusa Tenggara, South Kalimantan and West Kalimantan. Agarwood cultivation that is increasing especially after conducting several researches show that cultivated agarwood could provide feasible benefit (Marliani, 2008; Suharti, 2009; Tarmiji, 2009; The Angel, 2009) ^[71, 104, 108].

Agarwood exerts an important role in Indonesia as it contributes to country's foreign exchange. High economic value of agarwood has induced agarwood to become one of the prominent commodities in Indonesia (Pratiwi, 2010) ^[86]. Agarwood trees will not grow well on inundated soil, swamp area, soil solum thickness less than 50 cm, and quartz sand as well as soil with the acidity level less than 4 (Rizlani and Aswandi, 2009) ^[9]. Harvesting could be done before the trees die; however, agarwood is best harvested on the dead trees because three types of agarwood products: and ash/powder could be obtained all at once (Sumarna, 2007) ^[106].

Agarwood incenses have also been used as a fragrance in soaps and shampoos (Schippmann 2001) ^[96]. It is prescribed in traditional East Asian medicine to promote the flow of qi, relieve pain, arrest vomiting by warming the stomach, and to relieve asthma (Hajar, 2013) ^[45]. It is sold in the form of woodchips, wood pieces, powder, dust, oil, incense ingredients and perfume for several thousand US dollars per kilogram (Barden *et al.*, 2000, Gunn *et al.*, 2003 ^[15, 44], Compton, 2007) ^[31]. Agarwood chips start at US\$30 per kg up to \$9,000 per kg depending on its resin percentage (Babatunde, 2015) ^[13]. Agarwood oil fetches similarly high prices (Agarwood "Wood of Gods" International Conference 2003). The highest grade agar oil is produced in China and is reported to cost over Tk.0.9 million (US\$11,500) per kg. The second-best quality agar oil is sourced from Vietnam (Rahman *et al.*, 2015) ^[88].

A whole range of qualities and products are on the market, varying in quality with geographical location, botanical species, and the age of the specific tree, inoculation method and the section of the tree where the piece of agarwood stems from (Jung, 2011) ^[58]. The value of agarwood shipped out of Singapore alone each year has been estimated to exceed US\$1.2 billion (Hansen 2000) ^[46]. The current global market for agar oil and other related agarwood products is estimated to be in the range of US\$6 to 8 billion (Akter *et al.*, 2013) ^[5].

In Bangladesh, Aquilaria malaccensis (Alam, 2004), Aquilaria agallocha (Bhuiyan *et al.* 2009 and Rahman *et al.* 2015) ^[17, 88] are commonly cultivated. Agar trees were planted in monoculture and block plantations. 75% of the

plantations have been established on degraded land as well as denuded hills and rest 25% were established at the homesteads of farmers (Rahman *et al.*, 2015) ^[88].

After 3 to 5 years of nailing, the trees became ready for harvesting and the best time for harvesting is mid-October to mid-March (Chowdhury *et al.*, 2016 and Rahman *et al.*, 2015) ^[27, 88].

On an average, every enterprise uses 600 kg to 4 metric tons agarwood for producing 0.74-5.75 kg agar oil where production cost and total returns are BDT0.5 million and BDT1.3 million, respectively (Rahman *et al.*, 2015) ^[88] with a net profit of BDT0.8 million.

Abdin tried to find out some barriers toward the development of Bangladesh agarwood industry (Abdin 2014) ^[1]. Due to lack of contemporary price information of different markets, forest-based enterprises do not receive high prices for their products (Rahman *et al.*, 2009) ^[89].

Agarwood has been widely used as therapeutic perfumes, traditional medicine, religious purposes and aromatic food ingredient (Liu *et al.*, 2013) ^[67]. Some of the earliest known uses of agarwood were recorded in ancient literatures, religious scriptures and medical texts. The word “aloes” which means agarwood was found occurring in the Sanskrit poet, Kālidāsa that can be dated back to 4th-5th century (Lee and Mohamed, 2016) ^[65]. Meanwhile, the use of agarwood in the prescription of traditional Chinese medicine of the same period had also been recorded. The Chinese medicine uses it as a natural sedative, pain reliever, digestive aid and carminative (Ye *et al.*, 2016).

Agarwood has high demand throughout the world as a raw material for incense, perfume and medicine purposes, with Middle East and East Asia as the two major regions of consumption (Antonopoulou *et al.*, 2010) ^[8]. As the wealth of the consumer countries has gradually increased in the recent decades, the market’s demand for agarwood started to exceed its supply.

Global agarwood prices can be ranging from US\$ 20 – 6,000 per kilogram for the wood chips depending on its quality or US\$ 10,000 per kilogram for the wood itself (Abdin, 2014) ^[1]. In addition, the value of agarwood essential oil can be as high as US\$ 30,000 per kilogram. The annual global market for agarwood has been estimated to be in the range of US\$ 6 – 8 billion (Akter *et al.*, 2013) ^[5], yet a large number of the trades have not been recorded.

Aquilaria belongs to the Thymelaeaceae family of angiosperms, which is endemic to the Indomalayan realm. To date, there is a total of 21 Aquilaria species which have been documented and 13 of them are recognized as the agarwood-producing species (Lee and Mohamed, 2016) ^[65]. The destructive exploitation of agarwood, however, has badly affected the wild population of all Aquilaria species. As a consequence, the genus is now listed as endangered species and protected under Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) regulation due to a drastic declination of the species in the wild (Convention On International Trade In Endangered Species (CITES, 2004; Lee and Mohamed, 2016) ^[65, 29]. High demand of quality agarwood in conjunction with the depletion of the wild Aquilaria trees implied that the price of the agarwood will continue to soar. As an alternative, mass cultivation and large plantation of Aquilaria trees which serve as a sustainable source to obtain agarwood have greatly resolved the shortage of agarwood supply in the global market.

Since healthy Aquilaria tree does not form agarwood, leaving it worth next to nothing, the scarcity of naturally occurring agarwood has prompted the development of artificial agarwood-inducing methods. Efforts to artificially induce the agarwood formation can be traced back to as early as 300C.E. in the Chinese history, where it was recorded that resin deposition accompanied with color changes of internal tissues can happen within a year by injuring the trees (López-Sampson and Page, 2018) ^[68]. Besides mechanical wounding approach, the use of chemical, insect and pathogen-inducing techniques is increasingly common in the agarwood industry nowadays (Mohamed *et al.*, 2014; Kalita, 2015) ^[77]. All of these induction techniques in any case mimic the natural processes of agarwood formation, which have their own strengths and weaknesses. In this article, we endeavor to provide a more comprehensive coverage of existing induction methods and their development prospects using the advancement of biotechnology. To better understand the agarwood formation process, the molecular mechanism of secondary metabolite biosynthetic pathways underlying the resin production will also be elaborated. The indiscriminate harvesting of agarwood from natural habitats has seriously hampered natural regeneration of Aquilaria trees, thus threatening the survival of the species in the wild. In order to meet the high market demand yet to protect the species from extinction, mass plantations of Aquilaria trees have been established across the Asian countries to allow sustainable agarwood production (Azren *et al.*, 2018). Since agarwood formation in natural environment is a very long process which can take up to 10 years, the development of effective induction technology has received a great attention as it is extremely crucial to ensure the stability of agarwood yield from the domesticated Aquilaria trees.

Naturally, agarwood formation is often linked to the physical wounding or damage of Aquilaria trees caused by thunder strike, animal grazing, pest and disease infestations (Rasool and Mohamed, 2016; Wu *et al.*, 2017) ^[90, 115]. These events expose the inner part of the trees toward pathogenic microbes, which elicit the defense mechanism of Aquilaria to initiate the resin production. This natural formation process of agarwood has greatly inspired the development of diverse artificial induction methods (Table1). For example, many traditional induction approaches like nail in setting, holing, burning, trunk breaking and bark removal have adopted the concept of physically wound the trees (Mohamed *et al.*, 2010; Azren *et al.*, 2018). Although it is cost effective and requires only personnel with little or no scientific knowledge on agarwood, but these induction methods usually result in inferior quality and uncertain yield of agarwood.

With more understanding on Aquilaria-fungal interactions in promoting the agarwood formation, the induction methods gradually shifted from sole mechanical wounding into deliberate wounding coupled with the application of biological inoculums (Jong *et al.*, 2014) ^[57]. Many pure-culture strains of fungi isolated from natural agarwood were found to be effective biological agents to induce agarwood formation in healthy Aquilaria trees (Cui *et al.*, 2013; Sibirian *et al.*, 2015; Sangareswari *et al.*, 2016) ^[34, 99, 95]. The fungal infected Aquilaria trees were reported to deposit agarwood resin around the infected sites as barrier to prevent further fungal intrusion (Cui *et al.*, 2013; Rasool and Mohamed, 2016) ^[34, 90]. One obvious advantage of using

fungal inoculum is that it is generally believed to be safe for handling and eco-friendly. However, fungal inoculation will normally give rise to localized and inconsistent quality of agarwood due to the different fungal consortium used. As a solution, laborious holing process and long incubation time is required to maximize the colonized surface area on the tree to produce better quality of agarwood (Mohamed *et al.*, 2014) [77].

Aquilaria trees to initiate agarwood resin biosynthesis pathways (Liu *et al.*, 2013; Wu *et al.*, 2017) [67, 115]. Chemical inducers normally comprise of phytohormones, salts, minerals and biological derived substances (Zhang *et al.*, 2012; Liu *et al.*, 2013; Van *et al.*, 2015) [67, 123]. Besides, suitable delivery method is often developed together with the chemical formulations to ease the large-scale induction process, such as vessel equipped with transfusion needle (Yang *et al.*, 2014c). To date, several induction approaches have been developed based on the chemical induction concept such as cultivated agarwood kit (CA-kit), the whole-tree agarwood inducing technique (Agar-Wit) and biologically agarwood-inducing technique (Agar-bit). CA-kit is a combined method based on physical wounding and chemical induction, where the inducing agent is applied into the Aquilaria tree via an aeration device inserted into the wound (Blanchette and Heuveling, 2009) [21]. This method results in satisfying yield and quality, but the procedures are in some way conventional. On the other hands, Agar-Wit is a transpiration-assisted chemical treatment to form an overall wound in the tree, where the preloaded inducer in a transfusion set is distributed via plan transpiration (Liu Y. *et al.*, 2013) [67]. Through this method, a larger agarwood coverage area can be achieved, but unfortunately produces more decayed tissues. Similarly, Agar-bit method adopts the idea of distributing the inducing reagent by plant transpiration, except that the reagents are injected directly into the stems of the tree (Wu *et al.*, 2017).

Through chemical induction approach, the time-consuming holing process can be minimized as less induction sites are needed to deliver the inducers throughout the plants via transpiration process. Properly formulated inducer was shown to be able to produce artificial agarwood with quality closely resembled to those obtained from natural source (Liu *et al.*, 2013) [67]. In spite of the fast results and high yields, the application of chemical inducers still poses skepticism of toxicity on both human and environment. The currently adopted agarwood quality assessment in the market has been extensively reviewed by Liu *et al.* (2017). Recently, the metabolite analysis of agarwood has gained increasing attention as some studies showed that there is correlation of agarwood quality to its resin yield and metabolite constituents (Pasaribu *et al.*, 2015; Liu *et al.*, 2017). Many studies have been conducted to clarify the metabolite composition of agarwood obtained either from wild or artificially induced methods (Chen *et al.*, 2012; Gao X. *et al.*, 2014; Hashim *et al.*, 2014) [24, 47]. It was concluded that the composition of agarwood resin is mainly composed of the mixtures of sesquiterpenes and 2-(2-phenylethyl) chromones (PECs) (Naef, 2011; Chen *et al.*, 2012; Subasinghe and Hettiarachchi, 2015) [24, 78].

Meanwhile, the constituents of agarwood essential oil were shown primarily to be sesquiterpenoids (Fazila and Halim, 2012; Hashim *et al.*, 2014; Jayachandran *et al.*, 2014) [38, 47, 54]. Together, all of these major compounds and some low abundant volatile aromatic metabolites form the unique and

fragrant-smelling property of agarwood.

The number and types of agarwood metabolite constituents of each reported studies vary depending on the agarwood source, extraction methods and analysis approaches used (Fazila and Halim, 2012; Jong *et al.*, 2014; Pasaribu *et al.*, 2015) [38, 57]. Nonetheless, there are over 150 compounds as reviewed by Naef (2011) [78] have been identified thus far in agarwood from different sources.

The better insight of agarwood metabolites will definitely facilitate the identification of universally accepted biomarkers for agarwood grading. Since the publication of the comprehensive review of Naef (2011) [78] regarding the major constituents of agarwood, new compounds continue to be discovered in the later studies (Wu *et al.*, 2012a; Yang *et al.*, 2014b; Wang *et al.*, 2015). The number of discovered compounds in agarwood will certainly be further increased in the future.

Agarwood formation can be related to the self-defense mechanism of Aquilaria trees in response to (Gao *et al.*, 2012b; Singh and Sharma, 2015).

Due to the rampant destruction of natural habitats, most agarwood-bearing species have been relegated to the status of endangered species. In fact, all 19 known Aquilaria species are included under CITES (CITES, UNEP-WCMC Species Database: CITES-Listed Species, 2019) and the Red List of the IUCN (IUCN Red List of Threatened Species, 2019) [52].

In a recent study by Chen *et al.* (2018) [26], a fungus, *Rigidoporus vinctus*, was isolated from the inner layer of infected *A. sinensis* trees.

It was found that the Agar-Bit method up regulated the expression of some biological synthase genes (farnesyl diphosphate synthase, sesquiterpene synthase, and chalcone synthase) compared to the mechanically stimulated agarwood (Wu *et al.*, 2017; Tan *et al.*, 2019). Subsequently, Subasinghe *et al.* (2018) carried out an artificial induction by *Aspergillus niger* and *F. solanifer* agarwood resin formation in *Gyrinops wallatrees*.

Genetic engineering of plants has been used to produce terpenes by successfully expressing heterologous terpene synthase genes to produce novel monoterpenes, sesquiterpenes, and diterpenes (Besumbes *et al.*, 2004; Lucker *et al.*, 2004; Wu *et al.*, 2006) [114]. In one of the cited examples, Wu *et al.* (2006) [114] used the patchoulol synthase gene from patchouli to transform tobacco.

Trichome-specific promoters can be used to drive the accumulation of the terpenes specifically into leaf trichomes, which can confer ease of recovery of the compounds (Kim *et al.*, 2008; Shangguan *et al.*, 2008).

In a recent study, cloning of two new genes, AmSesTPS1 and AmGuaiS1, from *A. malaccensis* has been reported (Azzarina *et al.*, 2016).

Jasmonate induces expression of sesquiterpene synthase ASS1, which is a major enzyme in the biosynthesis of agarwood sesquiterpenes in *A. sinensis* (Xu *et al.*, 2017).

In a subsequent study, Kumeta and Ito (2010) further enhanced the understanding of agarwood biosynthetic machinery by revealing the genomic organization of the δ -guaiene synthase gene in *A. crassna*.

In a study by Kenmotsu *et al.* (2011), it was found that the biosynthesis of spirovetivane-type sesquiterpene in *A. microcarpais* triggered by the treatment with methyl jasmonate due to enhanced expression of farnesyl diphosphate synthase, calmodulin, and Rac/Rop GTPase

genes.

Tubulin, ribosomal protein, and glyceraldehyde 3-phosphate dehydrogenase genes were identified as the most stable reference genes for quantification of target gene expression in *A. sinensis* in the future (Gao *et al.*, 2012a).

Gao *et al.* (2012b) discovered 27 novel miRNAs in *A. sinensis*, and expression levels of 10 stress-responsive miRNAs were correlated with wounding. Eight of them were wound-responsive, which shows that the existence of miRNAs is indicative of their critical role in stress reactions, leading to agar formation (Gao *et al.*, 2012b).

References

1. Abdin MJ. The agar wood industry: yet to utilize in Bangladesh. *International Journal of Economics & Management Sciences* 2014;3(1):163-166. Doi: 10.2139/ssrn.2430055
2. Adelina N. *Aquilaria Malaccensis* Lam., SEED Leaflet, Forest & Landscape Denmark 2004.
3. Adijaya. Gaharu: Harta di kebun. trubus-online.co.id/mod.php?mod=publisher&p=allmedia&arti d=1625. Accessed 13 February 2009.
4. Akter N, Neelim AZ. Agarwood Plantation at BRAC Tea Estate: Introduction, Environmental Factors and Financial Analysis. Report Prepared for the BRAC Research and Evaluation Division 2008.
5. Akter S, Islam MT, Zulkefeli M, Khan SI. Agarwood production-a multidisciplinary field to be explored in Bangladesh. *International Journal of Pharmaceutical and Life Sciences* 2013;2:22-32. doi: 10.3329/ijpls.v2i1.15132
6. Alam SM. Production, processing and economic aspects of Agar tree (*Aquilaria malaccensis*) as an agroforestry species in Maulvibazar district of Bangladesh. M.Sc. thesis. Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh 2004.
7. Ali NS. Agarwood (Oud) 2006. Available at: <http://www.fragrantica.com/notes/Agarwood-Oud-114.html/>
8. Antonopoulou M, Compton J, Perry LS, Al-Mubarak R. The trade and use of agarwood (oudh) in the United Arab Emirates. Petaling Jaya, Selangor, Malaysia, TRAFFIC Southeast Asia 2010. http://www.traffic.org/species-reports/traffic_species_plants16.pdf
9. Aswandi. Budidaya Gaharu: Alternatif pemberdayaan masyarakat desa hutan. Accessed, 12 February 2009.
10. Azren PD, Lee SY, Emang D, Mohamed R. History and perspectives of induction technology for agarwood production from cultivated *Aquilaria* in Asia: a review. *J. For. Res* 2018;30:1-11. doi: 10.1007/s11676-018-0627-4.
11. Azzarina AB, Mohamed R, Lee SY, Nazre M. Temporal and spatial expression of terpene synthase genes associated with agarwood formation in *Aquilaria malaccensis* Lam. *N. Z. J. For* 2016;46:12. Doi: 10.1186/s40490-016-0068-9
12. Ba XD. A Vietnamese technique of agarwood inducement 2010. Available at: <http://www.agarwood.ning.com/profiles/blogs/agarwood-inducement/>
13. Babatunde OJ. Oud: Arabia's traditional scent. Available at: <http://www.masterpiece-n> Blanchette, R. A., and H. H. Van Beek. 2005. Cultivated agarwood. U.S. Patent No. 2015;6(848):211. In Washington, DC, U.S. g.com/ 2015/09/01/oud-arabias-traditional-scent/ Available at: <http://www.masterpiece-ng.com>.
14. Baksha MW, Akhter S, Basak AC, Rahman MS. Bangladesh agar chas o agar kutir silpo (Agar cultivation and agar cottage industry in Bangladesh). Bangladesh Forest Research Institute. Chittagong; (Bengali) 2009.
15. Barden A, Anak NA, Mulliken T, Song M. Heart of the Matter: Agarwood Use and Trade and Cites Implementation for *Aquilaria Malaccensis*. Traffic International. Cambridge, England. TRAFFIC Network Report 2000;46:17-18. https://www.traffic.org/forestry-reports/traffic_pub_forestry7.pdf
16. Besumbes O, Sauret-Güeto S, Phillips MA, Imperial S, Rodriguez-Concepción M, Boronat A. Metabolic engineering of isoprenoid biosynthesis in *Arabidopsis* for the production of taxadiene, the first committed precursor of Taxol. *Biotechnol. Bioeng* 2004;88:168-175. Doi: 10.1002/bit.20237
17. Bhuiyan MNI, Begum J, Bhuiyan MNH. Analysis of essential oil of eaglewood tree (*Aquilaria agallocha* Roxb.) by gas chromatography mass spectrometry. *Bangladesh Journal of Pharmacology* 2009;4:24-28.
18. Blanchette RA. Cultivated Agarwood -Training programs and Research in Papua New Guinea, Forest Pathology and Wood Microbiology Research Laboratory, Department of Plant Pathology, University of Minnesota 2006.
19. Blanchette RA. Successful Production of Cultivated Agarwood; A New Economy for Poor Rural People using Green Technology. Abstract. Second International Agarwood Conference 2007.
20. Blanchette R, Heuveling van Beek H. Cultivated Agarwood. US Patent 2005;6(848):211.
21. Blanchette R, Heuveling VBH. Cultivated Agarwood. U.S. Patent No. 7638145. Minnesota: University of Minnesota 2009.
22. Brechbill GO. The woody notes of fragrance. New Jersey, USA: Fragrance Book Inc 2012.
23. Chang YS, Nor Azah MA, Abu Said A, Lok EH, Reader S, Spiers A. Gaharu. FRIM Technical Information No 2002, 69.
24. Chen HQ, Wei JH, Yang JS, Zhang Z, Yang Y, Gao ZH. Chemical constituents of agarwood originating from the endemic genus *Aquilaria* plants. *Chem. Biodivers* 2012;9:236-250. Doi: 10.1002/cbdv.201100077.
25. Chen H, Yang Y, Xue J, Wei J, Zhang Z, Chen H. Comparison of Compositions and Antimicrobial Activities of Essential Oils from Chemically Stimulated Agarwood, Wild Agar wood and Healthy *Aquilaria Sinensis* (Lour.) Gilg trees. *Molecules* 2011;16:4884-4896.
26. Chen X, Liu Y, Yang Y, Feng J, Liu P, Sui C. Trunk surface agarwood-inducing technique with *Rigidoporus vinctus*: an efficient novel method for agarwood production. *PLoS One* 2018;13:e0198111. doi: 10.1371/journal.pone.0198111
27. Chowdhury M, Rahman A, Hussain MD, Kabir E. The economic benefit of agarwood production through aeration method into the *Aquilaria malaccensis* tree in Bangladesh. *Bangladesh Journal of Agricultural*

- Research 2017;42:191-196.
28. CITES. UNEP-WCMC Species Database: CITES Listed Species 2019. Species+, <https://www.speciesplus.net/#/elibrary> (Accessed July 02, 2019)
 29. CITES. The Trade and Use of Agarwood in Taiwan, Province of China. TRAFFIC East Asia-Taipei Available online 2005. <http://cites.org/sites/default/files/common/com/pc/15/X-PC15-07-Inf.pdf>
 30. CITES. Amendments to appendices I and II of CITES. In Proceedings of Thirteenth Meeting of the Conference of the Parties, Bangkok, Thailand, 2 October 2004.
 31. Compton J. Trade Matters: Regulating a Legal and Sustainable Agarwood Industry. 2nd International Agarwood Conference and Workshop. Bangkok and Koh Chang, Thailand 2007.
 32. Compton J, Ishihara A. The use and trade of agarwood in Japan. Southeast Asia and East Asia-Japan, TRAFFIC 2006. <http://144.76.93.178/sites/default/files/common/com/pc/15/X-PC15-06-Inf.pdf>
 33. Compton J, Ishihara A. The Use and Trade of Agarwood in Japan. TRAFFIC International, Cambridge 2004.
 34. Cui J, Guo S, Fu S, Xiao P, Wang M. Effects of inoculating fungi on agar wood formation in *Aquilaria sinensis*. Chin. Sci. Bull 2013;58:13280-3287. doi: 10.1007/s11434-013-5856-5.
 35. Dash MJK Patra, Panda PP. Phytochemical and anti microbial screening of extracts of *Aquilaria agallocha* Roxb. African Journal of Biotechnology 2008;7:3531-3534.
 36. Donovan D, Puri R. Learning from traditional knowledge of non-timber forest products: Penan Benalui and the autecology of *Aquilaria* in Indonesian Borneo. Ecology and Society 2004;9(3):3. [Online] URL:<http://www.ecologyandsociety.org/vol9/iss3/art>
 37. Espinoza EO, Lancaster CA, Kreitals NM, Hata M, Cody RB, Blanchette RA. Distinguishing wild from cultivated agarwood (*Aquilaria* spp.) using direct analysis in real time and time of-flight mass spectrometry. Rapid Communications in Mass Spectrometry 2014;28:1-9.
 38. Fazila KN, Halim KHK. Effects of soaking on yield and quality of agarwood oil. J Trop. For. Sci 2012;24:557-564.
 39. Fu JL. The Culture of Chinese Incense; QiLu Press: Ji'nan, China 2008, pp. 121-123.
 40. Gao X, Xie M, Liu S, Guo X, Chen X, Zhong Z. Chromatographic fingerprint analysis of metabolites in natural and artificial agarwood using gas chromatography-mass spectrometry combined with chemometric methods. J Chromatogr. B Analyt. Technol. Biomed. Life Sci 2014;967:264-273. Doi: 10.1016/j.jchromb.2014.07.039.
 41. Gao ZH, Wei JH, Yang Y, Zhang Z, Zhao WT. Selection and validation of reference genes for studying stress-related agarwood formation of *Aquilaria sinensis*. Plant Cell Rep 2012b;31:1759-1768. Doi: 10.1007/s00299-012-1289-x.
 42. Gao ZH, Wei JH, Yang Y, Zhang Z, Zhao WT. Selection and validation of reference genes for studying stress-related agarwood formation of *Aquilaria sinensis*. Plant Cell Rep 2012a;31:1759-1768. Doi: 10.1007/s00299-012-1289-x
 43. Gao ZH, Yang Y, Zhang Z, Zhao WT, Meng H, Jin Y. Profiling of microRNAs under wound treatment in *Aquilaria sinensis* to identify possible microRNAs involved in agarwood formation. Int. J Biol. Sci 2014;10:500-510. Doi: 10.7150/ijbs.8065.
 44. Gunn B, Stevens P, Singdan M, Sunari L, Chatterton P. Eaglewood in Papua New Guinea, Viewed on 15 March 2008. RMAP working paper no 51. Resource Management in Asia Pacific. First International Agarwood Conference TRP Vietnam. November 2003. http://rspas.anu.edu.au/papers/rmap/Wpapers/rmap_wp_51.pdf.
 45. Hajar A. Medical and other uses of Oudh, In Agarwood. The Most Expensive Wood Fragrance 2013.
 46. Hansen E. The hidden history of a scented wood. Saudi Aramco World 2000;51:1-13.
 47. Hashim Y, Ismail N, Abbas P. Analysis of chemical compounds of agarwood oil from different species by gas chromatography mass spectrometry (GCMS). IIUM Eng. J 2014;15:55-60. Doi: 10.31436/iiumej.v15i1.469.
 48. Hilton-Taylor C. IUCN Red List of Threatened Species. Gland, Switzerland, IUCN 2000.
 49. Hoang Ha Q, Nghi HT. Promoting agarwood-based Agroforestry systems in north central provinces of Viet Nam. Asia-Pacific Agroforestry Newsletter 2011;38(38):10-12.
 50. Islam MA, Hussain MD, Kabir E, Hasan MZ. Development of low cost agarwood essential oil extraction system for agar tree growers in Bangladesh. Journal of Bangladesh Agricultural University 2014;12(1):167-170.
 51. IUCN. IUCN Red List of Threatened Species. Available online: <http://www.iucnredlist.org> (accessed on 3 April 2009) 2008.
 52. IUCN. IUCN Red List of Threatened Species 2019. Available at: www.iucnredlist.org (Accessed March 27, 2019).
 53. IUCN. IUCN Red List of Threatened Species. Version 2013, 1. Available online: www.iucnredlist.org
 54. Jayachandran K, Sekar I, Parthiban KT, Amirtham D, Suresh KK. Analysis of different grades of agarwood (*Aquilaria malaccensis* Lamk.) oil through GC-MS. Indian J. Nat. Prod. Resour 2014;5:44-47.
 55. Jensen A. Valuation of Non-timber forest Products Value Chain. Forest Policy and Economics 2009;11:34-41.
 56. Jensen A. Species Monograph #12: *Aquilaria crassna*. Vientiane, Lao Tree Seed Project, Forest Research Centre, National Agriculture & Forestry Research Institute (available at LTSP office: 021-770074, ltsp1@laotel.com) 2002.
 57. Jong PL, Pascale T, Rozi M. Gas chromatography-mass spectrometry analysis of agarwood extracts from mature and juvenile *Aquilaria malaccensis*. Int. J. Agric. Biol 2014;16:644-648.
 58. Jung D. The Value of Agarwood. Reflections Upon Its Use and History in South Yemen. Universitätsbibliothel, Universität Heidelberg, 30 May 2011, (PDF) p. 4.
 59. Kalita J. Association of *Zeuzera conferta* Walker on agarwood formation in *Aquilaria malaccensis* Lamk. Asian J. Plant Sci. Res 2015;5:4-9.

60. Kenmotsu Y, Ogita S, Katoh Y, Yamamura Y, Takao Y, Tatsuo Y, *et al.* Methyl jasmonate-induced enhancement of expression activity of Am-FaPS-1, a putative farnesyl diphosphate synthase gene from *Aquilaria microcarpa*. *J. Nat. Med.* 2011;65:194-197. doi: 10.1007/s11418-010-0451-4
61. Kiet LK. History and ecology of agarwood in Vietnam. 1st International Agarwood Conference. November 10 – 15, Ho Chi Minh City, Vietnam 2003.
62. Kim SH, Chang YJ, Kim SU. Tissue specificity and developmental pattern of amorpho-4, 11-diene synthase (ADS) proved by ADS promoter-driven GUS expression in the heterologous plant, *Arabidopsis thaliana*. *Planta Med.* 2008;74:188-193. Doi: 10.1055/s-2008-1034276
63. Koczberski G, Curry GN, Gibson K. Improving Productivity of the Smallholder Oil Palm Sector in Papua New Guinea. RSPAS, Australian National University 2001;41:42-46.
64. Le CK. History and ecology of agarwood in Vietnam, Proceedings of the 1st International Agarwood Conference. The Tropical Rainforest Project 2003.
65. Lee SY, Mohamed R. The origin and domestication of *Aquilaria*, an important agarwood-producing genus, in *Agarwood: Science Behind the Fragrance*, ed. R. Mohamed. (Berlin: Springer Singapore) 2016, 1–20. Doi: 10.1007/978-981-10-0833-71
66. Lim TW, Anak. NA. Wood for the Trees: A Review of the Agarwood (Gaharu) Trade in Malaysia. TRAFFIC Southeast Asia. Petaling Jaya, Selangor, Malaysia 2010.
67. Liu Y, Chen H, Yang Y, Zhang Z, Wei J, Meng H, *et al.* Whole-tree agarwood-inducing technique: An efficient novel technique for producing high-quality agarwood in cultivated *Aquilaria sinensis* trees. *Molecules* 2013;18(3):3086-3106. Doi: 10.3390/molecules18033086.
68. López-Sampson A, Page T. History of use and trade of agarwood. *Econ. Bot.* 2018;72:107-129. Doi: 10.1007/s12231-018-9408-4.
69. Lucker J, Schwab W, Van Hautum B, Blaas J, Van der Plas L, Bouwmeester HJ, *et al.* Increased and altered fragrance of tobacco plants after metabolic engineering using three monoterpene synthases from lemon. *Plant Physiol* 2004;134:510-519. Doi: 10.1104/pp.103.030189
70. Mamat MF, Yacob MR, Fui LH, Rdam A. Costs and benefits analysis of *Aquilaria* species on plantation for agarwood production in Malaysia. *International Journal of Business and Social Science* 2010;1(2):162-174.
71. Marliani L. Suntikan inukolan - Sumber Majalah Trubus Indonesia 2009. <http://gaharuman.blogspot.com/2008/09/suntikan-inukolan-sumber-majalah-trubus.html>. Accessed 9 February 2009.
72. McKenna DJ, Hughes K. *The incense bible: Plant scents that transcend world culture, medicine, and spirituality.* New York: Routledge 2014.
73. Mei WL, Liu J, Li XN, Dai HF. Study on the chemical constituents from Chinese eaglewood in Hainan. *Journal of Tropical and Subtropical Botany* 2010;18:573-576.
74. Mei WL, Zeng YB, Wu J, Dai HF. Chemical composition and anti-MRSA activity of the essential oil from Chinese eaglewood. *Journal of Chinese Pharmaceutical Sciences* 2008;17:225-229.
75. Mohamed R, Lee SY. *Agarwood: Science behind the fragrance.* Singapore: Springer 2016a.
76. Mohamed R, Lee SY. Keeping up appearances: Agarwood grades and quality. In: *Agarwood: Science behind the fragrance*, ed. Rozi Mohamed, Singapore: Springer 2016b, 149-167.
77. Mohamed R, Jong PL, Kamziah AK. Fungal inoculation induces agarwood in young *Aquilaria malaccensis* trees in the nursery. *Journal of Forestry Research* 2014;25(1):201-204. Doi: 10.1007/s11676-013-0395-0.
78. Naef R. The volatile and semi-volatile constituents of agarwood, the infected heartwood of *Aquilaria* species: A review. *Flavour and Fragrance Journal* 2011;26:73-87.
79. Norsuzieana AL. Analysis volatile compound of gaharu oil composition via solid phase micro extraction (SPME). B. Sc. Thesis, Malaysia Pahang University, Malaysia 2009.
80. Page T, Awarau W. Performance of agarwood (*Aquilaria crassna*) seedling transplants improved by shade and fertilizer. *Forest Ecology and Management* 2012;265:258-269.
81. Paoli GD, Peart DR, Leighton M, Samsodin I. An ecological and economic assessment of the non timber forest product gaharu wood in Gunung Palung National Park, West Kalimantan, Indonesia. *Conservation Biology* 2001;15(6):1721-1732.
82. Pasaribu GT, Waluyo TK, Pari G. Analysis of chemical compounds distinguisher for agarwood qualities. *Indonesian J. For. Res* 2015;2:7. doi: 10.20886/ijfr.2015.2.1.1-7
83. Persoon GA. Agarwood: The Life of a Wounded Tree, Viewed on 15 March 2008. IIAS News Letter no. 45. Autumn 2007, pp. 24-25.
84. Persoon GA. Growing 'the wood of the gods': Agarwood production in Southeast Asia. In: *Smallholder tree growing for rural development and environmental services: Lessons from Asia*, vol. 5, eds. D. J. Snelder and R. D. Lasco 2008, 245-262, Dordrecht, Netherlands: Springer.
85. Pojanagaroon S, Kaewrak C. Mechanical methods to stimulate aloe wood formation in *Aquilaria crassna* Pierre ex H Lec (kritsana) trees. *ISHS Acta Hort* 2006;676:161-166.
86. Pratiwi ES, Turjaman M. Karakteristik lahan habitat pohon penghasil gaharu di beberapa hutan tanaman di Jawa Barat. *Info Hutan* 2010;8(2):129-139.
87. Qi SY, He ML, Lin LD, Zhang CH, Hu LJ, Zhang HZ. Production of 2-(2-phenylethyl) chromones in cell suspension cultures of *Aquilaria sinensis*. *Plant Cell, Tissue and Organ Culture* 2005;83(2):217-221.
88. Rahman M, Nath NM, Sarker S, Adnan M, Islam M. Management and Economic Aspects of Growing *Aquilaria agallocha* Roxb. in Bangladesh. *Small-scale forestry* 2015;14:459-478.
89. Rahman M, Rahman MM, Islam M. Financial viability and conservation role of betel leaf based agroforestry: an indigenous hill farming system of Khasia community in Bangladesh. *Journal of Forestry Research* 2009;20:131-136.
90. Rasool S, Mohamed R. Understanding agarwood formation and its challenges," in *Agarwood: Science*

- Behind the Fragrance, ed. R. Mohamed. (Berlin: Springer) 2016, 39–56. Doi: 10.1007/978-981-10-0833-7_3
91. Rhind JP. Fragrance and wellbeing: Plant aromatics and their influence on the psyche. London: Singing Dragon 2014.
 92. Rizlani C, Aswandi. Prospek budidaya gaharu secara ringkas. Accessed 18 February 2009.
 93. Rogers ZS.. A World Checklist of Thymelaeaceae (version 1). Missouri Botanical Garden: St. Louis, MO, USA 2009. Available online: <http://www.tropicos.org/project/thymelaeaceae> (accessed on 9 July 2009).
 94. Roy K, Samal NR, Roy MB, Mazumdar A. Soil Carbon and Nutrient Accumulation under Forest Plantations in Jharkhand state of India. Clean - Soil, Air, Water 2010;38:706-712.
 95. Sangareswari M, Parthiban KT, Kanna SU, Karthiba L, Saravanakumar D. Fungal microbes associated with agarwood formation. American Journal of Plant Sciences 2016;7:1445-1452. doi: 10.4236/ajps.2016.710138
 96. Schippmann U. Medicinal Plants Significant Trade Study: CITES Project S-109, Plants Committee Document, and PC 99.1. 3 (rev.). BfN 2001.
 97. Shangguan X, Xu B, Yu Z, Wang L, Che X. Promoter of a cotton fibre MYB gene functional in trichomes of Arabidopsis and glandular trichomes of tobacco. J. Exp. Bot 2008;59:3533-3542. doi: 10.1093/jxb/ern204
 98. Sherbinin A, Wey LKV, Mc Sweeney K, Aggarwal R, Barbieri A, Henry S, *et al.* Rural Household Demographics, Livelihoods and the Environment. Global Environmental Change 2008;18:38-53.
 99. Sibirian RH, Siregar UJ, Siregar IZ, Santoso E. Identification of morphological characters of *Aquilaria microcarpa* in the interaction with *Fusarium solani*. Int. J. Sci. Basic Appl. Res 2015;20:119-128.
 100. Singh B, Sharma RA. Plant terpenes: defense responses, phylogenetic analysis, regulation and clinical applications. 3 Biotech. 2015;5:129-151. doi: 10.1007/s13205-014-0220-2.
 101. Soehartono T, Newton AC. Conservation and sustainable use of tropical trees in the genus *Aquilaria*. I. Status and distribution in Indonesia. Biol. Conse 2000;96:83-94.
 102. Subasinghe SMCUP, Hettiarachchi DS. Characterisation of agarwood type resin of *Gyrinops walla* Gaertn growing in selected populations in Sri Lanka. Ind. Crops Prod 2015;69:76-79. Doi: 10.1016/j.indcrop.2015.01.060.
 103. Subasinghe SMCUP, Hitihamu HID, Fernando KMPEP. Use of two fungal species to induce agarwood resin formation in *Gyrinops walla*. J. For. Res 2018;30:721-726. Doi: 10.1007/s11676-018-0654-1
 104. Suharti S. Prospek pengusahaan gaharu melalui pola pengelolaan hutan berbasis masyarakat (PHBM). Prosiding Workshop: Pengembangan Teknologi Produksi Gaharu Berbasis Pemberdayaan Masyarakat Sekitar Hutan. Pusat Penelitian dan Pengembangan Hutan dan Konservasi Alam, Bogor Bekerjasama dengan ITTOPD 425/06 Rev 2009;1(1):219 pp.
 105. Sumarna Y. The Effects of Seed Maturity Condition and Media Type on Growth of Agarwood Seedlings of Karas Trees (*Aquilaria malaccensis* Lamk.), Journal Penelitian Hutan dan Konservasi Alam 2008;5(2):129-135.
 106. Sumarna Y. Budidaya dan rekayasa produksi gaharu. Temu Pakar Pengembangan Gaharu. Direktorat Jenderal RLPS, Jakarta 2007, 15p.
 107. Tan CS, Isa NM, Ismail I, Zainal Z. Agarwood induction: current developments and future perspectives. Front. Plant Sci 2019;10:122. Doi: 10.3389/fpls.2019.00122
 108. Tarmiji M. Perhitungan kelayakan usaha gaharu. Accessed 14th November 2009.
 109. The Angel. Siapkan masa depan, ayo tanam gaharu <http://The.angel>. Accessed 28th March 2009.
 110. Uddin MS, Mukul SA, Khan MASA, Alamgir M, Harun MY, Alam MS. Small-Scale Agar (*Aquilaria agallocha* Roxb.) Based Cottage Enterprises in Maulvibazar District of Bangladesh: Production, Marketing and Potential Contribution to Rural Development. Small-Scale Forestry 2008;7(2):139-149.
 111. Van T, Van Do L, Son T NH, Sato T, Kozan O. Impacts of biological, chemical and mechanical treatments on sesquiterpene content in stems of planted *Aquilaria crassna* trees. Agroforest. Syst 2015;89:973-981. Doi: 10.1007/s10457-015-9829-3.
 112. Wu B, Lee J, Lim C, Dong Jia S, Won Kwon S, Seo Hwang G. Sesquiterpenoids and 2-(2-phenylethyl)-4H-chromen-4-one (=2-(2-phenylethyl)-4H-1-benzopyran-4-one) derivatives from *Aquilaria malaccensis* agarwood. Helv. Chim. Acta 2012a;95:636-642. Doi: 10.1002/hlca.201100409
 113. Wu B, Won Kwon S, Seo Hwang G, Hill Park J. Eight new 2-(2-phenylethyl)chromone (= 2-(2-Phenylethyl)-4H-1-benzopyran-4-one) derivatives from *Aquilaria malaccensis* agarwood. Helv. Chim. Acta 2012b;95:1657-1665. Doi: 10.1002/hlca.201200069.
 114. Wu S, Schalk M, Clark A, Miles RB, Coates RC. Redirection of cytosolic or plastidic isoprenoid precursors elevates terpene production in plants. Nat. Biotechnol. 2006;24:1441-1447. Doi: 10.1038/nbt1251
 115. Wu ZQ, Liu S, Li JF, Li MC, Du HF, Qi LK. Analysis of gene expression and quality of agarwood using Agarbit in *Aquilaria sinensis*. J Trop. For. Sci 2017;29:380-388. doi: 10.26525/jtfs2017.29.3.380388
 116. Wyn LT, Anak NA. Wood for the trees: A review of the agarwood (gaharu) trade in Malaysia. Petaling Jaya, Selangor, Malaysia: TRAFFIC Southeast Asia 2010.
 117. Xu YH, Liao YC, Lv FF, Zhang Z, Sun PW, Gao ZH. Transcription factor AsMYC₂ controls the jasmonate-responsive expression of ASS1 regulating sesquiterpene biosynthesis in *Aquilaria sinensis* (Lour). Gilg. Plant Cell Physiol 2017;58:1924-1933. Doi: 10.1093/pcp/pcx122
 118. Yang DL, Wang H, Guo ZK, Li W, Mei WL, Dai HF, *et al.* Fragrant agarofuran and eremophilane sesquiterpenes in agarwood 'Qi-Nan' from *Aquilaria sinensis*. Phytochem. Lett 2014b;8:121-125. Doi: 10.1016/j.phytol.2014.03.003.
 119. Yang M, Fu H, Liang Y, Huang H, Zhao B, Xie C. Modified transfusion devices, inducer, and procedure for agarwood-inducing by infusion technique. J Chem. Pharm. Res 2014c;6:2566-2571.
 120. Ye W, Wu H, He X, Wang L, Zhang W, Li H. Transcriptome sequencing of chemically induced *Aquilaria sinensis* to identify genes related to agarwood

- formation. *PLoS One* 2016;11:e0155505.
doi: 10.1371/journal.pone.0155505.
121. Zeng D, Mao R, Li L. Restoration Effects of Young Plantations on Magnesite Mine Spoil in North China. International Conference on Bioinformatics and Biomedical Engineering. 18-20 January 2010. Chinese Academy of Science, Shenyang China 2010, 1-5.
 122. Zhang L, Brockelman WY, Allen MA. Matrix analysis to evaluate sustainability: The tropical tree *Aquilaria crassna*, a heavily poached source of agarwood. *Biological Conservation* 2008;141(6):1676-1686.
 123. Zhang XL, Liu YY, Wei JH, Yang Y, Zhang Z, Huang JQ. Production of high-quality agarwood in *Aquilaria sinensis* trees via whole-tree agarwood-induction technology. *Chin. Chem. Lett* 2012;23:727-730. Doi: 10.1016/j.ccllet.2012.04.019.
 124. Zich F, Compton J. The Final Frontier: Towards Sustainable Management of Papua New Guinea's Agarwood Resource. A TRAFFIC Oceania Report in conjunction with WWF South Pacific Programme 2002.
 125. Zich F, Compton J. The Final Frontier. Towards sustainable management of Papua New Guinea's agarwood resource. Kuala Lumpur. TRAFFIC Oceania, Sydney 2001.