



E-ISSN: 2663-1067  
P-ISSN: 2663-1075  
IJHFS 2021; 3(2): 11-15  
Received: 18-05-2021  
Accepted: 22-06-2021

**Ogori AF**

Department of Home Science and Management, Faculty of Agriculture, Federal University, Gashua, P.M.B.1005 Gashua, Yobe State, Nigeria

**Amove J**

Department of Home Science and Management, Faculty of Agriculture, Federal University, Gashua, P.M.B.1005 Gashua, Yobe State, Nigeria

**Evi-Parker P**

Department of Home Science and Management, Faculty of Agriculture, Federal University, Gashua, P.M.B.1005 Gashua, Yobe State, Nigeria

**Corresponding Author:**

**Ogori AF**

Department of Home Science and Management, Faculty of Agriculture, Federal University, Gashua, P.M.B.1005 Gashua, Yobe State, Nigeria

## Quality characteristics of jam from pineapple, cucumber and Jathropa

**Lilian Blanco and Thippi Thiagarajan**

**Abstract**

Production of jams is one of the preserving techniques to extend whole fruit utilization. Fruits are constituents and important sources of bioactive components, therefore their mixed Jams from various fruits, can be a proper solution against degenerative chronic diseases. In the present study, physicochemical and sensory properties of mixed jam from pineapple fruit, Jathropa leaves and cucumber fruit were evaluated. Mixed jam were prepared by sorting, washing, and blending different ratios of apple, cucumber and jathropa (T1(85:10:5), T2 (80:10:10), T3 (75:10:15), and T4(70:10:20) respectively. The obtained results on proximate includes moisture content, protein values, ash content, fat content, crude fiber and carbohydrate values which ranged from 3.83-15.61% moisture content, 0.06-0.707% protein, 0.014-0.110% Ash, 0.023-0.145% fat, 0.123-0.592% crude fibre and 83.25-96.14% carbohydrate value respectively. The minerals contents includes calcium, zinc, magnesium, potassium, sodium, and iron and ranged from 1.87-2.87mg for Ca, 0.03-0.10mg for Zn, 0.11-0.7mg for Mg, 0.12-0.61mg for K, 1.78-2.59mg for Na and 0.10-0.67mg for Fe respectively. Vitamin C and Beta carotene were evaluated and reported thus; 691.35-436.95mg/100g and 0.73-1.44mg/100g respectively. The results from sensory attributes and physicochemical properties reveal rich mixed jam product from pineapple fruits, jathropa leaves and cucumber fruit. T4, sample ratio portrayed the most acceptable jam product.

**Keywords:** Mixed jam, fruit, vegetable, blends, physicochemicals, sensorial, quality

**Introduction**

Prepared fruits, vegetables and sugar often canned or sealed for long-term storage are referred to as fruit preserves, Jam, a products from a whole fruit, cut in pieces or crushed. The fruit is heated with water and sugar for pectin activation within the fruit. The mixture is then put into the containers. The fruit preserves often involves the addition of pectin as a gelling agent, sugar or honey may be used as well (Afoakwa, 2006) <sup>[1]</sup>.

*Jatropha tanjorensis* popular in Mexico and originated in Central America. According to Egbon *et al.* (2013) <sup>[4]</sup>, it can be grown on waste land and in local palace. The liquid extracted from it leave is drunk to quickly beef up blood volume in the body. It can also be used as alternative to lettuce in salad preparation. Olayiwola *et al.* (2004) <sup>[10]</sup> opined that the leaf of *Jatropha tanjorensis* is been used locally and also consumed as food vegetable. It is popular as a natural remedy against diabetes for rural dweller in Nigeria.

Cucumber is the fruit of *Cucumis sativus* plant. Cucumbers are high in water and low in calories, fat, cholesterol, and sodium (George, 2007) <sup>[6]</sup>. They are noted for their mild and soothing taste with a high water content. Pineapples (*Ananas comosus*) are tropical fruit that are rich in vitamins, enzymes and antioxidants. Despite their sweetness, pineapples are low in calories. The fruit is actually made of many individual berries that fuse together around a central core. It is a non-toxic compound which has a number of potential therapeutic applications, including treatment of trauma, inflammation, auto immune disease, enhancement of immune response and malign (Bhaskar, 2011) <sup>[3]</sup>. Jams made from a combination of fruits and vegetables are rare. This work is aimed at producing a jam with fruits and vegetables complimenting each other resulting in higher nutritional quality, as well as an appealing, sensory and synergistic bioactive values.

**Scientific hypothesis**

Mixed fruit jam from pineapple, jathropa and cucumber were produced as home preserves. The following hypothesis were tested: either that jathropa and cucumber blend ratios had any effect on the chemical, vitamin, mineral and sensorial properties or not on jam preserve. Increasing or decreasing the blend ration percentage of jathropa and cucumber on pineapple derived mixed jam may improve the physicochemical properties (protein, crude

fibre, ash fat) values of mixed jam preserve. Increasing or decreasing the percentage blend ratio of jathropha and cucumber in proportional ration in making jam may improve the vitamin and mineral contents of mixed fruit jam preserve. Increasing the or decreasing the blend ration percentage may improve the sensorial properties (mouthfeel, appearance, and general acceptability) of pineapple derived mixed jam

## Material and Methodology

### Materials

Mature and healthy pineapple and cucumber fruits were purchased from North Bank market Makurdi and selected on the basis of shape size, uniformity, color and damages. Any fruits with the injuries and disease were discarded. Fresh leaves of jathropha also were collected from the environment housing the College of Food Technology and human Ecology complex of the University of Agriculture Makurdi.

### Fruit and Vegetable Preparations

Selected fruits and vegetables are sorted, washed to remove debris and adhering soils and dirt from their skin. In the next step, fruits were peeled and cut into small pieces the crushed/blended. The pulp from the three individual fruits and vegetable were extracted in the same manner to obtain three pulps. The pulp and sugar were mixed and heated. The soluble solid formations were monitored during the process until the brix of 55°C was attained (Kawther *et al.*, 2013).

To prepare mixed Jam, pectin was used as solution base. These solutions were prepared by the addition of pectin to hot water and heating to reach a homogeneous solution mixture. The pectin solution, the fruit pulp, sugar and citric acid were added at this stage. The jam was ready when the bubbles formed at the sides of the vessels were observed. The hot jam was poured in clean, dry, wide-mouthed jars or bottles and cooled to 35 °C in a water bath for gel formation and reduction of thermal stress. The final jam was then stored at ambient temperature for 24 H.

### Physico-Chemical Analysis of pineapple, cucumber and Jathropha blended Jam

The moisture contents, pH, ash, fat, protein and crude fibre contents of the jam was determined in accordance with AOAC (2012) [2] method.

#### Determination of pH

The pH was standardized and the electrode was rinsed with distilled water, the electrode was then dipped into 5g of the sample which was dissolved in 50 mL of water.

#### Determination of Moisture Content

Two gram (2g) of the prepared samples were weighed in petri dishes and covered immediately, then transferred into an oven, uncovered and heated at 130-150°C for 3 H. After the heating the samples were removed, placed in a desiccator and allowed to cool for 15 min before weighing. This was repeated until constant weights were recorded. The loss in weight from the original (before heating) was reported as the percentage moisture content equation 1).

$$\text{Eq (1): Moisture Content} = \frac{\text{Weight loss} \times 100\%}{\text{Weight of sample}}$$

#### Determination of Protein Content

AOAC (2012) [2] method was adopted. 0.8g of the prepared samples was digested in kjedahl digestion system under a fume chamber. The digesta was allowed to cool and then distilled into boric acid containing bromocresol green indicators after it is appropriately diluted first with water, followed by sodium thiosulphate and sodium hydroxide solutions. In the next step, the samples were titrated against 0.1N Hydrochloric acid (HCL) solutions. Blank titrations were similarly carried out and the percentage protein content was calculated by equation 2.

$$\text{Eq (2): \%Nitrogen} \times 6.25(1\text{ml of } 0.1\text{N HCL} = 0.0014\text{gN})$$

$$\text{Nitrogen} = \frac{\text{Titre value} - \text{blank} \times 0.0014\text{N} \times 100\% \times 25}{\text{Weight of sample} \times 5\text{ml aliquot}}$$

#### Determination of Ash Content

AOAC (2012) [2] method was adopted. Five grams of the prepared sample was weighed in duplicates into ash dishes that have been previously weighed. The dishes were placed into a muffle furnace and ignited at 550 ± 10 °C for 5 H, cooled and then weighed to constant weight. The resulting ash was calculated based on equation 3.

$$\text{Eq (3): Ash Content} = \frac{\text{Weight of ash} \times 100\%}{\text{Weight of sample}}$$

#### Determination of Fat Content

AOAC (2012) [2] method was adopted. Two (2g) of the prepared sample was weighed into soxhlet thimbles and fixed into extraction flask of given weight. Extraction was carried out using diethyl ether for 5 H. At the completion, the diethyl ether was removed by evaporation on an electrical bath and the remaining fat in the flask dried at 60 °C for 30 min in oven, then cooled for 15 min and weighed. Equation 4 was used to evaluate fat content.

$$\text{Eq (4): Ash Content} = \frac{\text{Weight of fat} \times 100\%}{\text{Weight of sample}}$$

#### Determination of Crude Fibre Content

The AOAC (2012) [2] method was adopted. One gram of the sample was weighed. Then 100 mL of Trichloroacetic acid (TCA) digesting reagent was added. The solution was brought to boil and reflux for exactly 40min at 50-60°C. The flask was removed from the heater, cooled a little and the solution was filtered through Whatman filter paper. The residue was washed with hot water and with methylated spirit. The filtrate was transferred to a muffle furnace and ignited at 550°C for 30 min, cooled and weighed. The percentage crude fibre content was calculated as:

$$\% \text{ Crude fibre} = \frac{\text{the loss in weight after incineration}}{\text{original weight}} \times 100$$

#### Determination of Carbohydrate Content

Carbohydrate value of sample was determined in accordance with the method described by AOAC (2012) [2]. Equation was used to calculate carbohydrate percent.

$$\text{Carbohydrate percent} = 100 - (\% \text{ Moisture} + \% \text{ Fat} + \% \text{ Protein} + \% \text{ Crude fibre} + \% \text{ Ash})$$

#### Total Titratable Acid

Ten (10g) of the sample was dissolved in 100 mL of distilled

water. 10 mL aliquot of the supernatant was titrated with 0.1N NaOH and phenolphthalein as an indicator. Titrable acidity was reported based on citric acid following the below equation.

% citric acid = volume of NaOH used  $\times$  0.1N  $\times$  ml equivalent of citric acid  $\times$  100

#### Determination of Total Soluble Solids (TSS)

Dry empty dishes were weighed and 5g of the samples were added. These were placed on a boiling water bath and left to stand until the water was evaporated from the samples. The samples were then placed in oven at 102 °C for 2.5 H. This was allowed to cool and readings taken. Equation was used to calculate the SS percent.

$$\% \text{ total solid} = \frac{\text{Weight of residue} \times 100}{\text{Weight of sample}}$$

#### Determination of Total Sugar Content (°Brix)

AOAC (2012) [2] method was adopted. Hand held sugar refractometer instrument was used. The prism of the refractometer was cleaned and a drop of the sample was placed on the prism and closed. The total sugar Content (°Brix) was read off the scale of the refractometer when held close to the eyes.

#### Determination of Minerals

The mineral elements such as Calcium, Zinc, Magnesium, Potassium, Sodium and Iron were determined using the method by AOAC (2012) [2]. One gram of each sample was weighed into 100 mL round bottom flask, and 5 mL of perchloric acid was added and heated over an electric heater in a fume chamber until the solution becomes colourless. Each of the solution was made up to 10 mL mark with distilled water and the diluted sample set aside for further studies. The Calcium, Zinc, Magnesium, Potassium, Sodium and Iron content were analyzed using the Atomic Absorption Spectrophotometer.

#### Determination of Vitamins

The Vitamin A ( $\beta$ -carotene) and Vitamin C (ascorbic acid) was determined using the method by AOAC (2012) [2]. Ten grams of each sample was weighed into a 250mL flask and 50mL acetone was added. It was allowed to stand for 2 H with occasional shaking, and then filtered. The filtrate was measured and equal volume of saturated NaCl was added to wash the filtrate (carotene extract). The resulting mixture was shaken and transferred to a separating funnel to remove the layer of the extracted carotene. The supernatant was washed again with equal volume of 100% Potassium Trioxocarbonate (IV) ( $K_2CO_3$ ), which separates and ultimately washed with 10-20 mL of distilled water. Water carotene was separated and the carotenoid extracted. The absorbance was read in a spectrophotometer at 326 nm wavelength using 50:50 acetone and low boiling petroleum ether solution as the blank.

#### Sensory Evaluation

Sensory evaluation was carried out to assess consumers' preference using a 9-point hedonic scale where 1 represented dislike extremely while 9 represented like extremely. Ten panelists were employed to evaluate the attributes of appearance or colour, odour, taste and general

acceptability of the jam. The panelists were selected out of students of the Department of Food Science and Technology, University of Agriculture, Makurdi. For each sample, 10g of jam was served in white disposable plates with some slice of bread. The plates were coded with three digit random numbers undistributed randomly among panelists. The evaluation took place at 10a.m. in the sensory evaluation laboratory.

#### Statistical Analysis

Result data were expressed as the mean values and standard deviation of two determinations. Data were analysed using a one-way analyses of variance (ANOVA) using Statistical Package for Social Science (SPSS) version 20.0 software, 2011 ( $p < 0.05$ ) (Lawless and Heymann, 2010) [4].

#### Results and Discussion

Table 2 depicted the results of the proximate composition of mixed fruit and vegetable jam prepared from pineapple, cucumber and jathropa. Moisture contents of various samples differed significantly ( $p < 0.05$ ). The least and the most amounts of moisture content were control sample 3.82 moisture content is least and sample 75:10:15 (15.61%) had the highest value respectively. The shelf life of the control sample (100% pineapple) might be stable compared to those from other samples. However, the spreadability of control (100% pineapple) was low due to lower moisture content observed.

Protein increased significantly from 0.061%  $\pm$  0.001 to 0.707%  $\pm$  0.007, where control sample and sample 70:10:20 depicted the least and the most amounts respectively. This increase arose from the addition of cucumber and Jathropa leaves. Similarly, it was found that control sample and sample 75:10:15 had the least and the most amounts ash contents. Higher amounts of ash in sample 75:10:15 could be due to the addition of cucumber and jathropa leaves which makes mixed jam a good source of minerals as observed by Egbon *et al.* (2013) [4]. Considering the ash content results shown in Table 2, control sample A and sample 75:10:15 differ significantly at ( $p < 0.05$ ), while other samples do not vary significantly from each other at ( $p < 0.05$ ).

The fat content of the mixed fruit and vegetable jam samples increased from 0.023% (control sample) to 0.145% in sample (85:10:05) and subsequently decreased. Sample 85:10:05 has the highest value (0.145%), while control sample has the lowest value (0.023%). The increase in the fat content was due to the addition of cucumber, which depicted a considerable level of fat as described by Uzuazokaro *et al.* (2018), whereas, the subsequent decrease was owing to increasing addition of Jathropa. The result also showed no significant differences between other samples at ( $p < 0.05$ ). Crude fiber of the mixed fruit and vegetable jam samples was shown to increase from 0.123 – 0.592 %. Addition of jathropa and 10 % cucumber to the blend might be the reason for increasing crude fiber.

Carbohydrate content of the jam decreased from 96.14% (control sample) to 83.25% in sample (75:10:15). The varying values of the carbohydrate (calculated by difference) is due to uneven heat treatment given to the sample, hence an increase in the carbohydrate value of sample 70:10:20 (87.58%) was reported. Results showed a significant difference ( $p < 0.05$ ) in the carbohydrate content of samples.



The mineral and vitamin contents of different blends of mixed fruit and vegetable jam prepared from pineapple, cucumber and jathropha are shown in Table 3. Calcium, Zinc, Magnesium, Potassium and Iron contents of the mixed fruit and vegetable jams were higher than control at ( $p < 0.05$ ).

The addition of cucumber and Jathropha was reported to increase the Calcium, Zinc, Magnesium, Potassium and Iron values of the mixed jam. Deficiency of zinc had been shown to aggravate carbohydrate intolerance in certain individuals (Franco *et al.*, 2000) [13]. Also, an increase in calcium intake had been reported beneficial and likely to reduce osteoporosis and diabetes in older people surviving from diabetes (Iwalewa *et al.*, 2005) [9].

The contents of  $\beta$ -carotene and vitamin C differed from each other at ( $p < 0.05$ ). This could become increased in the percentage addition of Jathropha leaves.

Table 4 show results from physiochemical analysis. The low and high values of TTA were reported in sample 70:10:20 (9.15%) and sample 75:10:15 (16.70%) respectively. The decrease in the TTA might be due to the increase of jathropha and addition of 10% cucumber in samples 85:10:05 and sample 70:10:20. The pH of the jam increased significantly ( $p > 0.05$ ). The samples became more alkaline because of the increase in the addition level of blended jathropha and the use of 10% blended cucumber in samples 85:10:05 to sample 70:10:20. The alkaline nature of this jam

product could be an advantage against aerosol viruses (WHO, 2020). No significant difference reported in decrease of pH from control sample to sample 80:10:10, whereas the increase in pH was significantly different from samples 80:10:10 to 70:10:20

The brix values of the samples decreased at ( $p > 0.05$ ). This reduction was due to the increase in the addition level of blended jathropha and the use of 10% blended cucumber in samples 85:10:05 to sample 70:10:20.

Total soluble solid values ranged from 84.38–96.17. It increased from control Sample with the low value to sample 70:10:20 with the high value. It increased with an increase in minerals resulting from addition level of blended jathropha and the use of 10% blended cucumber.

The results of sensory evaluation tests have been summarized in Table 5. The organoleptic evaluation remains the final judge of the food quality. The appearance of the jam samples was affected by the blended fruits and vegetables. The taste for the control was ranked significantly ( $p < 0.05$ ) higher than sample 70:10:20, however shown to be the least preferred with lowest score among samples. This is probably due to the high amount of Jathropha in sample 70:10:20. The spread ability of the control sample is significantly different from the others due to high and uneven heat treatments. The overall acceptability of the samples was not significantly different at ( $p < 0.05$ ).

**Table 1:** Product Formulation of mixed fruit and vegetable jam from Pineapple, Cucumber and Jathropha.

Samples	Pineapple	Cucumber	Jathropha	Total
A(control)	100	0	0	100
B	85	10	5	100
C	80	10	10	100
D	75	10	15	100
E	70	10	20	100

A = 100% Pineapple, B = 85% Pineapple; 10% Cucumber; 5% Jathropha, C = 80% Pineapple; 10% Cucumber; 10% Jathropha, D = 75% Pineapple; 10% Cucumber; 15% Jathropha, E = 70% Pineapple; 10% Cucumber; 20% Jathropha

**Table 2:** The proximate composition of jam from pineapple, cucumber and Jathropha

Samples PS:CF:JF	Proximate Composition (%)					
	Moisture	Protein	Ash	Fat	Fibre	CHO
100:0:0	3.82 <sup>a</sup> ±0.03	0.061 <sup>c</sup> ±0.001	0.014 <sup>c</sup> ±0.001	0.023 <sup>d</sup> ±0.004	0.123 <sup>c</sup> ±0.001	96.14 <sup>a</sup> ±0.001
85:10:05	10.17 <sup>d</sup> ±0.02	0.134 <sup>d</sup> ±0.016	0.039 <sup>b</sup> ±0.0001	0.145 <sup>a</sup> ±0.004	0.217 <sup>d</sup> ±0.009	89.29 <sup>b</sup> ±0.009
80:10:10	15.20 <sup>b</sup> ±0.02	0.405 <sup>c</sup> ±0.008	0.041 <sup>b</sup> ±0.0001	0.136 <sup>ab</sup> ±0.001	0.305 <sup>c</sup> ±0.008	83.90 <sup>d</sup> ±0.008
75:10:15	15.61 <sup>a</sup> ±0.02	0.506 <sup>b</sup> ±0.008	0.080 <sup>ab</sup> ±0.002	0.134 <sup>bc</sup> ±0.001	0.417 <sup>b</sup> ±0.007	83.25 <sup>e</sup> ±0.007
70:10:20	10.88 <sup>c</sup> ±0.02	0.707 <sup>a</sup> ±0.007	0.110 <sup>a</sup> ±0.001	0.124 <sup>c</sup> ±0.006	0.592 <sup>a</sup> ±0.007	87.58 <sup>c</sup> ±0.003
LSD	0.08	0.02	0.003	0.010	0.016	0.257

Values are mean and standard deviations of two replications, Values within a column with same superscript are not significantly different ( $p > 0.05$ ), Key: PS = Pineapple pulp; CS = Cucumber pulp; JF = Blended Jathropha leaves pulp, LSD = Least significant difference

**Table 3:** The vitamins and mineral composition of jam from pineapple, cucumber and Jathropha

Samples PS:CF:JF	Vitamin C (g/kg)	$\beta$ -Carotene (mg/100g)	Ca (mg/kg)	Zn (mg/kg)	Mg (mg/kg)	K (mg/kg)	Na (mg/kg)	Fe (mg/kg)
100:0:0	691.35 <sup>a</sup> ±0.49	1.44 <sup>a</sup> ±0.02	0.12 <sup>c</sup> ±0.004	1.87 <sup>c</sup> ±0.027	0.03 <sup>d</sup> ±0.002	0.11 <sup>c</sup> ±0.001	1.78 <sup>b</sup> ±0.001	0.10 <sup>c</sup> ±0.008
85:10:05	680.65 <sup>b</sup> ±0.21	1.25 <sup>b</sup> ±0.04	0.34 <sup>d</sup> ±0.003	2.00 <sup>d</sup> ±0.004	0.06 <sup>c</sup> ±0.001	0.34 <sup>d</sup> ±0.001	1.79 <sup>b</sup> ±0.003	0.20 <sup>d</sup> ±0.001
80:10:10	569.35 <sup>c</sup> ±0.49	1.12 <sup>c</sup> ±0.02	0.40 <sup>c</sup> ±0.001	2.20 <sup>c</sup> ±0.003	0.07 <sup>b</sup> ±0.002	0.45 <sup>c</sup> ±0.002	1.81 <sup>b</sup> ±0.017	0.33 <sup>c</sup> ±0.013
75:10:15	477.45 <sup>d</sup> ±0.21	0.81 <sup>d</sup> ±0.01	0.55 <sup>b</sup> ±0.001	2.71 <sup>b</sup> ±0.017	0.09 <sup>a</sup> ±0.001	0.65 <sup>b</sup> ±0.001	1.87 <sup>b</sup> ±0.024	0.45 <sup>b</sup> ±0.011
70:10:20	436.95 <sup>e</sup> ±0.07	0.73 <sup>e</sup> ±0.01	0.61 <sup>a</sup> ±0.016	2.87 <sup>a</sup> ±0.001	0.10 <sup>a</sup> ±0.002	0.71 <sup>a</sup> ±0.008	2.59 <sup>a</sup> ±0.140	0.67 <sup>a</sup> ±0.029
LSD	0.887	0.056	0.019	0.037	0.004	0.010	0.163	0.041

Values are mean and standard deviations of two replications, Values within a column with same superscript are not significantly different ( $p > 0.05$ ), Key: PS = Pineapple pulp; CS = Cucumber pulp; JF = Blended Jathropha leaves pulp, LSD = Least significant difference

**Table 4:** The Physico-chemical property of jam from pineapple, cucumber and Jathropha

Samples	Physico-chemical Properties			
	TTA (%)	pH	Brix (°brix)	TSS (%)
PS:CF:JF				
100:0:0	0.1670 <sup>a</sup> ±0.98	3.16 <sup>c</sup> ±0.09	79.05 <sup>a</sup> ±0.07	84.38 <sup>c</sup> ±0.02
85:10:05	0.1270 <sup>b</sup> ±0.28	3.18 <sup>c</sup> ±0.01	73.15 <sup>b</sup> ±0.07	84.79 <sup>c</sup> ±0.02
80:10:10	0.1225 <sup>b</sup> ±0.35	3.23 <sup>c</sup> ±0.03	69.00 <sup>c</sup> ±0.00	87.31 <sup>bc</sup> ±3.59
75:10:15	0.0925 <sup>c</sup> ±0.35	3.41 <sup>b</sup> ±0.01	66.20 <sup>d</sup> ±0.14	89.11 <sup>b</sup> ±0.02
70:10:20	0.0915 <sup>c</sup> ±0.21	3.85 <sup>a</sup> ±0.07	64.10 <sup>e</sup> ±0.14	96.17 <sup>a</sup> ±0.04
LSD	0.0134	0.14	0.26	4.12

Values are mean and standard deviations of two replications, Values within a column with same superscript are not significantly different ( $p>0.05$ ), Key: PS = Pineapple pulp; CS = Cucumber pulp; JF = Blended Jathropha leaves pulp; LSD = Least significant difference

**Table 5:** The Sensory properties of jam from pineapple, cucumber and Jathropha

Samples	Sensory Properties				
	Appearance	Taste	Aroma	Spread-ability	General Acceptability
PS:CF:JF					
100:0:0	8.20 <sup>a</sup>	7.93 <sup>a</sup>	7.67 <sup>a</sup>	3.87 <sup>b</sup>	7.27 <sup>a</sup>
85:10:05	7.07 <sup>b</sup>	7.93 <sup>a</sup>	7.67 <sup>a</sup>	7.27 <sup>a</sup>	7.73 <sup>a</sup>
80:10:10	6.47 <sup>bc</sup>	7.93 <sup>a</sup>	7.27 <sup>a</sup>	7.67 <sup>a</sup>	7.60 <sup>a</sup>
75:10:15	5.73 <sup>cd</sup>	7.33 <sup>a</sup>	6.93 <sup>ab</sup>	7.80 <sup>a</sup>	7.53 <sup>a</sup>
70:10:20	5.07 <sup>d</sup>	7.07 <sup>a</sup>	6.47 <sup>b</sup>	7.60 <sup>a</sup>	6.87 <sup>a</sup>
LSD	1.05	0.89	0.70	1.05	0.83

Values with same superscript within a column are not significantly different at ( $p>0.05$ ), Key: PS = Pineapple pulp; CS = Cucumber pulp; JF = Blended Jathropha leaves pulp, LSD = Least significant difference

### Conclusions

The addition of cucumber and Jathropha leaf to pineapple for jam production increased the nutrient constituents of the jam. Jam containing 70% pineapple, 10% cucumber and 20% Jathropha leaves pulp blends were most acceptable. Based on the result of the study, production of jam from 20% Jathropha leaves, 10% cucumber blends and 70% pineapple pulp are recommended.

### References

- Afoakwa E, Oguseye O, Annor G, Ashong J. Optimization of the processing conditions and quality characteristics of watermelon jams using response surface methodology, paper presented at the 13th world food congress, 2006, 221-222. <https://doi.org/10.7287/PEERJ.PREPRINTS.1777V1>.
- AOAC. Official methods of Analysis of Association of Official Analytical Chemistry International, 19th edition. Washington, DC, 2012.
- Bhaskar JJ. Banana (*Musa sp. varelakki bale*) Flower and Pseudostem: Dietary Fiber and Associated Antioxidant Capacity. *Journal of Food Chemistry Biotechnology* 2011;4:223-228. <https://doi.org/10.1021/jf204539v>.
- Egbon EE, Ize-Iyamu OK, Okojie VU, Egbon IE. Proximate and mineral composition of jathrophatanjorensis. *Chemical and Process Engineering Research* 2013;17(1):21-23.
- Amit Kumar. Utilization of bioactive components present in pineapple waste: A review. *Pharma Innovation*. 2021;10(5):954-961.
- George M. *The World's Healthiest Foods*, 2nd edition. George Mateljan Foundation, Washington 2007, 288-289.
- Lawless HT, Heymann H. *Sensory Evaluation of Food Principles and Practices* (2nd Edition), Springer, New York, 2010.
- Kawther MYK, Abdel MS, Zakaria AS. Extraction of pectin from lemon and orange fruit peels and its utilization in Jam making. *International Journal of Food Science and Nutrition Engineering* 2013;3(5):81-84. <https://doi.org/10.5923/j.food.20130305.01>.
- Iwalewa EO, Adewunmi CO, Omisore NO, Adebajji OA, Azike CK. Pro-antioxidant effects and cytoprotective potentials of nine edible vegetables in Southwest Nigeria. *Journal of Medicinal Food* 2005;8:539-544. <https://doi.org/10.1089/jmf.2005.8.539>.
- Olayiwola G, Iwalewa EO, Omobuwajo OR, Adeniyi AA, Verspohi EJ. The antidiabetic potential of jathrophatanjorensis leaves. *Nigerian journal of natural products and medicine* 2004;8:55-58. <https://doi.org/10.4314/njnpm.v8i1.11817>.
- Uzuazokaro MA, Okwesili FCN, Chioma AA. Phytochemical and proximate composition of cucumber (*Cucumis sativus*) fruit from Nsukka, Nigeria. *African Journal of Biotechnology*, 2018;17(38):1215. <https://doi.org/10.5897/AJB2018.16410>.
- WHO. 2019 Coronavirus pandemic, issues hygiene and safety. <https://www.who.int/emergencies/disease/novel-coronavirus-2019/media-resources/news>
- Franco R, Alejandra C, Blanca O, Murillo OS, Sergio LB, Paula V, *et al.* Evaluation of chemical composition and antibacterial activity of essential oils of turmeric 2007;4:208-220.